

CHAPTER 4

COASTAL WETLANDS

Coordinating Lead Authors

Hilary A. Kennedy (UK), Daniel M. Alongi (Australia), Ansarul Karim (Bangladesh)

Lead Authors

Guangcheng Chen (China), Gail L. Chmura (Canada), Stephen Crooks (USA), James G. Kairo (Kenya), Baowen Liao (China), Guanghui Lin (China)

Contributing Author

Tiffany G. Troxler (USA)

Review Editors

Nuria Marba Bordalba (Spain) and Georgi Karl Hiebaum (Bulgaria)

Contents

4.1	Introduction	4.5
4.1.1	Land remaining in a land-use category.....	4.9
4.1.2	Conversion from a land-use category that includes coastal wetlands.....	4.12
4.1.3	Conversion to a land-use category that includes coastal wetlands	4.13
4.2	Land remaining in a land-use category	4.13
4.2.1	Biomass	4.14
4.2.2	Dead Organic Matter	4.22
4.2.3	Soil Carbon.....	4.25
4.2.4	Non-CO ₂ emissions	4.35
4.3	Conversion from a land-use category that includes coastal wetlands	4.41
4.3.1	CO ₂ emissions and removals	4.42
4.3.2	Non-CO ₂ emissions	4.42
4.4	Conversion to a land-use category that includes coastal wetlands.....	4.42
4.4.1	CO ₂ emissions and removals	4.42
4.4.2	Non-CO ₂ emissions	4.42
4.5	Completeness, Times Series consistency, Quality Assurance and Quality Control.....	4.43
4.5.1	Completeness	4.43
4.5.2	Developing a consistent time series.....	4.43
4.5.3	Quality assurance and quality control	4.43
4.5.4	Reporting and documentation.....	4.44
4.6	Future Methodological Development	4.44
References	4.45
Annex 4	4.66

Equations

Equation 4.1	Annual change in carbon pools in land remaining in a land use category	4.13
Equation 4.2	Change in soil C in coastal wetlands with aquaculture, salt production activities in three phase of sub-activities and extraction	4.27
Equation 4.3	Annual change in carbon stocks in drained organic and mineral soils	4.32
Equation 4.4	Annual change in soil C in restored and created coastal wetlands.....	4.33
Equation 4.5	Direct N ₂ O emissions from Aquaculture use	4.38

Figures

Figure 4.1	Decision tree to estimate greenhouse gas emissions and removals due to management activities in coastal wetlands.....	4.7
------------	---	-----

Tables

Table 4.1	Management Activities in Coastal Wetland Ecosystems Related to Land-Use Category and Chapter section	4.5
Table 4.2	Management Activities, Tier 1 Equations or Default EF for Biomass Pool Changes	4.14
Table 4.3	Carbon content of aboveground mangrove forest biomass (% DW)	4.17
Table 4.4	Aboveground Biomass in Mangrove forests (tonnes DW ha ⁻¹)	4.17
Table 4.5	Aboveground biomass growth in mangrove forests (tonnes DW ha ⁻¹ yr ⁻¹).....	4.18
Table 4.6	Ratio of belowground biomass to aboveground biomass (R) in mangroves forests	4.18
Table 4.7	Wood Density (D) of Common Mangrove Tree Species.....	4.18
Table 4.8	Aboveground and Belowground biomass for Tidal Marshes (tonnes C ha ⁻¹).....	4.19
Table 4.9	Aboveground biomass for seagrass meadow	4.20
Table 4.10	Ratio of belowground biomass to aboveground biomass (R) for seagrass meadow	4.20
Table 4.11	Carbon content of aboveground and belowground seagrass biomass (% DW)	4.21
Table 4.12	Management Activities, Tier 1 Equations and default EF for Dead Organic Matter C Pool Changes	4.22
Table 4.13	Tier 1 default values for litter and dead wood carbon stocks.....	4.24
Table 4.14	Management Activities, Tier 1 Approaches and Tier 1 Equations for Soil Carbon Pool Changes	4.26
Table 4.15	Emission factors (EF) associated with construction (EFCONSTR) of aquaculture (AQ), salt production (SP) and extraction (EXT) on organic soils (tonnes C ha ⁻¹) at start of activity	4.29

Second Order Draft

86	Table 4.16	Emission factors (EF) associated with construction (EF_{CONSTR}) of aquaculture (AQ), salt production (SP) and extraction (EXT) on mineral soils (tonnes C ha^{-1}) at start of activity	4.29
87			
88	Table 4.17	Annual emission factors (EF) associated with use (EF_{USE}) of aquaculture (AQ) on organic and mineral soils ($\text{tonnes C ha}^{-1} \text{ yr}^{-1}$)	4.30
89			
90	Table 4.18	Annual emission factors (EF) associated with abandonment after aquaculture (AQ) or salt production (SP) under saturated conditions (EF_{ABAN-S}) on organic and mineral soils and harvesting of aquatic resources (EF_{HARV}) in mangrove forests ($\text{tonnes C ha}^{-1} \text{ yr}^{-1}$)	4.30
91			
92			
93	Table 4.19	Annual emission factors (EF) associated with nutrient enrichment (EF_{NUTR}) on organic and mineral soils ($\text{tonnes C ha}^{-1} \text{ yr}^{-1}$)	4.31
94			
95	Table 4.20	Annual emission factors (EF) associated drainage (EF_{DR}) on organic and mineral soils ($\text{tonnes C ha}^{-1} \text{ yr}^{-1}$)	4.32
96			
97	Table 4.21	Annual emission factors (EF) associated with restoration or creation ($EF_{RES/CRE}$) on organic and mineral soils ($\text{tonnes C ha}^{-1} \text{ yr}^{-1}$) after 20 years of vegetation reestablishment	4.34
98			
99	Table 4.22	Activities and Tier 1 Assumptions for non- CO_2 emissions for the Ecosystems Affected	4.36
100	Table 4.23	Emission factors for N_2O emission from aquaculture in coastal wetlands	4.38
101	Table 4.24	Emission factors for CH_4 emissions from nutrient enrichment (agricultural run-off and aquaculture effluent) ($\text{tonnes CH}_4 \text{ ha}^{-1} \text{ yr}^{-1}$)	4.39
102			
103	Table 4.25	Emissions factors for N_2O emissions from nutrient enrichment (aquaculture effluent) ($\text{kg N}_2\text{O ha}^{-1} \text{ yr}^{-1}$)	4.39
104			
105	Table 4.26	Emission factors for CH_4 from unmanaged coastal wetlands for Tier 1 estimation of rewetting and restoration	4.40
106			
107			

Boxes

108			
109	Box 4.1	The following represent examples of different management activities that do or do not result in a change in land-use category	4.8
110			
111			
112			
113			
114			
115			
116			
117			
118			
119			
120			
121			
122			
123			

4.1 INTRODUCTION

This chapter provides guidance on estimating and reporting anthropogenic greenhouse gas (GHG) emissions and removals from managed coastal wetlands.

Coastal wetlands are defined here as organic and mineral soils vegetated by vascular plants that are covered or saturated for all or part of the year by tidal freshwater or salt water (>0.5 ppt) at or near the coast. The boundary of coastal wetlands is recognized as the landward extent of tidal inundation and extending seaward to the maximum depth of vascular plant vegetation with 95% of seagrass meadows being found shallower than 40 meters water depth (Duarte 1991). This description is derived from recent definitions in a specialized treatise on coastal wetlands (Perillo et al. 2009) and from a global treatise on all wetlands (Aber et al. 2012). This definition and this chapter thus refer specifically to tidal freshwater and salt marshes, seagrass meadows, and mangrove forests. All vegetation types occur on both mineral and organic soils. The guidance in this chapter applies to all managed coastal wetlands that can occur in any IPCC land category and addresses all IPCC pools (Table 4.1).

Supplementary guidance (Table 4.1) is structured to address greenhouse gas emissions and removals for Land Remaining in a Land-use category (Section 4.2), Land Converted to another Land-use category in which the land that is converted is a coastal wetland (Section 4.3), and Conversion to a Land-Use category that includes coastal wetlands (Section 4.4) regardless of how it is classified. Management activities that have significant impacts on national level CO₂ emissions and removals vary depending on country circumstances. This chapter provides guidance on management activities that will not be relevant for all countries but is meant to be inclusive of those activities that are common in managed coastal wetlands. The guidance is thus structured to aid the inventory compiler to identify areas of managed coastal wetlands and report on management activities that impact greenhouse gas emissions and removals. It should be noted also that depending on how managed coastal wetlands are classified, the effect of the management activity may or may not result in a land-use change. This is an especially important consideration for the guidance provided in this chapter. Inventory compilers should also be mindful that, *significant* greenhouse gas emissions and removals can result in an IPCC Wetlands land category that is *key* regardless of whether a land-use conversion has occurred (in reference to *significant* and *key*, see Chapter 4, Volume 1 of the *2006 IPCC Guidelines*). As such, to ensure complete coverage, this guidance enables the use of the IPCC Land category reporting system. Methodological assumptions and issues more specific to the three categories are discussed in the corresponding sections of this chapter. Readers are referred to Volume 4 of the *2006 IPCC Guidelines* for many of the basic equations to estimate greenhouse gas emissions, but methodological assumptions and other deviations are highlighted throughout the text, especially with regard to new activities including aquaculture and salt production where new methodological guidance is provided. The Decision Tree guides the inventory compiler to the appropriate Tier level estimation methodology as is followed in the *2006 IPCC Guidelines* (Figure 4.1).

TABLE 4.1 MANAGEMENT ACTIVITIES IN COASTAL WETLAND ECOSYSTEMS RELATED TO LAND-USE CATEGORY AND CHAPTER SECTION

LAND REMAINING IN A LAND-USE CATEGORY: Section 4.2 (This section covers management activities that occur in coastal wetland ecosystems and that may or may not result in a conversion to another land-use category.)		
Activity	Ecosystems Affected	Subtype
Aquaculture (AQ)	Mangrove forests, Tidal marshes	Fish, shrimp ponds
	Seagrass meadows	Fish cages, pens
Salt Production (SP)	Mangrove forests, Tidal marshes	
	Seagrass meadows	NA ¹
Extraction (EXT)	Mangrove forests, Tidal marshes, Seagrass meadows	Dredging; Port, harbour and marina construction; filling
Harvesting of Aquatic Resources (HARV)	Mangrove forests	Wood, Charcoal, Land clearing
	Tidal marshes	Grazing (cattle, sheep)
	Seagrass meadows	Fish, Shellfish

Second Order Draft

Nutrient Enrichment (NUTR)	Mangrove forests, Tidal marshes, Seagrass meadows	
Hydrologic/Sediment Diversion (DIV)	Mangrove forests, Tidal marshes	Impoundments, barriers, oil and gas extraction - resulting in soil elevation loss
	Seagrass meadows	NA
Drainage (DR)	Mangrove forests, Tidal marshes	Agriculture, forestry, mosquito control
	Seagrass meadows	NA
Restoration (RES) and Creation (CRE)	Mangrove forests, Tidal marshes	Reestablishment of vegetation following hydrologic or sediment modifications
	Seagrass meadows	Reestablishment of vegetation following water quality improvements
Other Activities	Mangrove forests, Tidal marshes	Recreation (hunting), fire management
	Seagrass meadows	Recreation (anchoring, mooring, boating), salt production
CONVERSION FROM A LAND-USE CATEGORY THAT INCLUDES COASTAL WETLANDS Section 4.3 (This section covers the conversion of a Land-use category that includes coastal wetlands to another Land-use Category, primarily including (1) drainage in which the soil water table is lowered for conversion to Cropland or Grassland or (2) excavation of soil resulting in conversion to Settlement.		
CONVERSION TO A LAND-USE CATEGORY THAT INCLUDES COASTAL WETLANDS Section 4.4 (This section covers restoration and creation of coastal wetland ecosystems in which a Land-use conversion occurs.		
NA =not applicable; ¹ no available Tier 1 guidance		

WHAT IS NOT COVERED IN THIS CHAPTER


For constructed wetlands that occur in coastal zones that are modified to receive and treat waste water, refer to Chapter 6 (this Supplement). Chapter 6 also covers semi-natural treatment wetlands which are natural wetlands where wastewater has been directed for treatment but the wetland is otherwise unmodified. With reference to nutrient enrichment as an activity in coastal wetlands, this chapter on coastal wetlands covers CO₂, CH₄ and N₂O emissions from aquaculture and effluent derived from aquaculture as well as CH₄ emissions from untreated, uncollected agricultural runoff and leaching resulting from direct nutrient application to terrestrial soils. Nutrient enrichment effects on CO₂ emissions from this type of agricultural run-off affecting mangroves and tidal marshes is not covered as there is inadequate information to provide emission factors for all pools and risk of double-counting across pools. Direct and indirect N₂O emissions from soils associated with N applied to terrestrial soils (direct and indirect) are covered in Chapter 11, Volume 4 of the *2006 IPCC Guidelines*. This Chapter only covers N₂O emissions during stocking of aquaculture and as a result of effluent from aquaculture in the coastal wetland. This chapter covers emissions from these lands, but excludes carbon losses or gains via tidal exchange (see Section 4.6).

SPECIAL CONSIDERATIONS FOR THE APPLICATION OF THE WETLANDS SUPPLEMENT GUIDANCE FOR COASTAL WETLANDS

Special considerations for the application of the Wetlands Supplement guidance for coastal wetlands include an understanding of: A) how to identify coastal wetlands, B) treatment of managed lands in coastal wetlands, C) identification of coastal ecosystems, D) soil type, and other generally applied assumption for soils.

A. Identification of coastal wetlands

Figure 4.1 Decision tree to estimate greenhouse gas emissions and removals due to management activities in coastal wetlands.



*A key source/sink category is defined in Chapter 4, Volume 1 of the 2006 IPCC Guidelines, "as one that is prioritised within the national inventory system because its estimate has a significant influence on a country's total inventory of greenhouse gases in terms of the absolute level, the trend, or the uncertainty in emissions and removals". The 2006 IPCC Guidelines recommend that the key category analysis is performed at the level of land remaining in or converted to a land-use category. If CO₂, CH₄ or N₂O emissions/removals from coastal wetlands are subcategories to a key category, these subcategories should be considered as significant if they individually accounts for 25-30% of emissions/removals for the overall key category (see Figures 1.2 and 1.3 in Chapter 1, Volume 4 of the 2006 IPCC Guidelines).

Second Order Draft

The first priority for inventory compilers is to determine if the managed wetlands under consideration conform to the definition and boundary of coastal wetlands that have been described. In many regions, these boundaries have been mapped using aerial photographs or remotely sensed data, and confirmed by field inspections, and in the form of freely available data (refer to respective sections on Activity Data and Annex 4.2). Wetlands are by definition lands with minimal slope and vegetated by species adapted to temporary, regular or permanent coverage by tidal freshwater or salt water (>0.5ppt) and saturated soil conditions. Many plant species are salt tolerant or salt resistant and may be adapted to grow fully submerged. Thus, wetlands can be identified by the presence of these specialized plant species or by soil characteristics that reflect flooded conditions (e.g., organic carbon content >12% or in mineral soils the presence of mottling).

B. Treatment of managed lands in coastal wetlands

The second priority for inventory compilers is to apply the concept of managed land in order to focus time and resources and report on management activities that impact anthropogenic emissions and removals of greenhouse gases at the national level. The activities that will result in the largest greenhouse gas emissions or removals are those that: 1) include removal and oxidation of soils (eg. aquaculture, salt production, extraction, drainage), 2) have both initial and sustained effects (eg. aquaculture, salt production, restoration) or 3) affect more than one pool. Additionally, for example, those activities which result in removal of forest biomass would have a larger impact than those that result in removal of tidal marsh biomass. So, the land area in which the activity occurs (coastal wetland ecosystem type) is also an important consideration in documentation and reporting of management activity impacts. Furthermore, IPCC *good practice* guidance requires documentation and reporting of all IPCC pools.

C. Identification of ecosystem types

A further consideration when applying the guidance in this chapter is the ecosystem type that is the subject of the management activity. Just as all management activities will not occur in all countries, not all management activities occur in all ecosystem types nor will all coastal wetland ecosystem types be considered managed. Coastal wetlands can occur in any IPCC land-use category regardless of how the land is classified (see examples Box 4.1). While countries should follow their own national definitions of coastal wetland ecosystem types, some general features that may help in consistent identification can be found throughout this guidance.

BOX 4.1. THE FOLLOWING REPRESENT EXAMPLES OF DIFFERENT MANAGEMENT PRACTICE THAT DO OR DO NOT RESULT IN A CHANGE OF A LAND-USE CATEGORY.

For Land remaining in a land-use category:

When areas of fish pens constructed within seagrass meadows are classified as Wetlands.

When tidal marshes or mangrove forests are classified as Wetlands and exploited for aquaculture or drained for agriculture that is also classified as Wetlands.

When selective harvesting or forest clearing occurs in mangrove forests that are classified as Wetlands to provide wood for fuel, charcoal or construction.

For Conversion from a land-use category that includes coastal wetlands:

When areas of fish pens constructed within seagrass meadows are classified as Settlements or Other Land.

When tidal marshes or mangrove forests are classified as Wetlands and exploited for aquaculture or drained for agriculture with that land area classified as Cropland. If the soils are organic and permanently drained for Cropland uses, refer to guidance in Chapter 2 of this Supplement. If the soils are mineral and permanently drained for Cropland uses, refer to guidance in Chapter 5 of the 2006 IPCC Guidelines.

When mangrove forests classified as Forest Land are deforested, drained and maintained as grazing land and subsequently classified as Grassland.

D. Identification of soil types

A unique aspect of this chapter in the Supplement is that coastal wetland ecosystems can occur with either mineral or organic soils. For some management activities, soils may be aggregated and the same emission factor applied regardless of the soil type (i.e. drainage, restoration and nutrient enrichment). However, emission factors do vary based on ecosystem types and Tier 1 methods disaggregate ecosystem types on this basis. For other management activities, emission factors are disaggregated by soil type (i.e. aquaculture, salt production). This

refinement results from the differing methodologies applied in these cases. Once management activities are identified, the inventory compiler should consider whether soil type must be determined. Thus, the fourth priority for inventory compilers is to distinguish organic from mineral soils in coastal wetlands once management activities and ecosystem types have been identified for Tier 1 estimation in the absence of soil map data. There are two assumptions that can be applied at Tier 1:

- (i) Assumption 1: Soils which are or become colonised by seagrass meadows are assumed to be mineral.
- (ii) Assumption 2: Soils in or at the mouth of estuaries or adjacent to any river characterised by a large and/or mountainous catchment and high flow are assumed to be mineral soils. For all other mangrove forests (sometimes termed oceanic mangroves) and tidal marshes the soils are assumed to be organic. See Durr et al. 2011 for additional national level guidance.

4.1.1 Land remaining in a land-use category

The known management activities that occur in coastal wetlands and that may not result in their conversion to another IPCC land-use category depending on how those lands are classified are aquaculture, salt production, extraction, harvesting of aquatic resources, nutrient enrichment, hydrologic/sediment diversion, drainage, restoration/creation and other activities (Table 4.1). In the following sub-section, these activities and how they affect the coastal wetlands they primarily impact are described. All methodological guidance is found in section 4.2. When management activities do not result in a land-use category conversion, follow guidance in section 4.2. When a land-use category conversion is involved, refer to guidance in sections 4.3 and 4.4 which direct the inventory compiler to the appropriate methodological guidance.

Aquaculture (shrimp ponds, fish ponds, fish cages)

There exists a range of aquaculture practices, but the most important, and ones that commonly occur in coastal wetlands, are fish farming and shrimp ponds (World Bank 2006).

Mangrove Forests and Tidal Marshes. Shrimp and fish ponds in these settings are identified through three phases: construction, use and abandonment. In the first phase they are constructed by clearing vegetation, levelling, followed by excavation of surface soils to build containing berms within which water is held. During construction, carbon from the cleared biomass and excavated soils is lost from storage and subsequently oxidized (World Bank 2006). While in use (second phase), ponds are stocked with fish and shrimp, and soils underlying ponds receive excess organic matter leading to an enhancement of benthic CO₂ fluxes from the oxidation of the organic matter (Burford & Langton 2001). N₂O will be emitted from aquaculture systems primarily as a by-product of the conversion of ammonia (contained in fish urea) to nitrate through nitrification and nitrate to N₂ gas through denitrification (Hu et al., 2012). While CH₄ production may occur if bottom muds are anoxic, some of the CH₄ will be oxidized as it passes through the water column, which must be oxygenated for fish culturing (Bose et al., 1991). However, there is considerable uncertainty regarding N₂O and CH₄ fluxes associated with aquaculture and additional research is necessary. When stocking of the ponds is discontinued, the ponds are abandoned. On abandonment, the area utilized for aquaculture is often left with saturated soils. In this guidance, this Chapter covers CO₂ emissions and removals from all phases of aquaculture and N₂O emissions only during the period when the ponds are being stocked.

Seagrass Meadows. Fish farming is a rapidly expanding industry in Asia (FAO 2007a). The fish are retained in floating cages or pens made up of nets often fastened to bamboo sticks that are anchored in the sediment (Alongi et al., 2009). Fish farming generates organic waste that impacts the immediate environment resulting in nutrient enrichment of the water column and organic enrichment of sediments beneath the farm. Nutrient over-enrichment of seagrass meadows has led to water and sediment quality deterioration, algal overgrowth, and loss of seagrass (Apostolaki et al., 2010; Apostolaki et al., 2011). Increased organic matter accumulation increases benthic metabolism leading to anoxic conditions and high sediment sulfide concentrations have been associated with seagrass die-off events (Holmer & Heilskov 2000; Borum et al., 2005). Gas bubbles have been observed in the sediments underlying fish pens and the sulphate concentration were depleted in the surface layers, thus it is thus likely that methane production occurs in these settings but no measurements have been made to date (Holmer et al., 2002).

Salt production

Mangrove Forests and Tidal Marshes. The Ramsar convention recognizes salt exploitation sites or solar salterns as a type of wetland. These are sites where salt is produced by evaporating seawater. The area covered can be

Second Order Draft

extensive and sites are found along tropical and subtropical coasts worldwide (Oren 2009). Many solar salterns have long histories, on the order of centuries (Thiery and Puente 2002). More recently, salt production has been shown to displace mangroves on the coast of Java (Sukardjo 1993). Some production sites, as in San Francisco Bay and on the Camargue, have been abandoned and restoration activities are underway. Establishment of the production system is similar to the construction phase of aquaculture and can include clearing vegetation, levelling, diking and constructing channels to develop a multi-pond production system (Oren 2002). Scientific evidence is not adequately developed to provide default emission factors for effects on CO₂ emissions while the salt production ponds are in use. The CO₂ emissions and removals of salt production are similar as those of aquaculture during construction and abandonment and so similar guidance is given in both instances.

Extraction (dredging; port, harbour and marina construction; filling)

Mangrove Forests and Tidal Marshes. Dredging excavates soil often leading to loss of coastal wetlands. In mangrove forests and tidal marshes this most often occurs by direct physical removal of biomass and soil. The CO₂ emissions and removals associated with this extraction are the same as those of aquaculture during construction and so the same guidance is given in both instances. The area that has had the soil removed is sometimes subsequently filled with other soil, whether extracted within or outside the coastal ecosystem. Filling of coastal wetlands is also considered a significant management activity affecting CO₂ emissions; if filling occurs, it is preceded by extraction. Similarly as for extraction, CO₂ emissions and removals must be reported. Once the extraction is complete, the area that has had the soil removed may become covered with water and CO₂ emissions are assumed to be negligible.

Seagrass Meadows. Extraction effects on seagrass meadows can take two forms: 1) direct via on-site physical removal of soil and biomass (Zainal et al. 1993, Ertlinmeyer et al 2012) or 2) indirect via sediment resuspension that causes high turbidity and reduces water quality impacting biomass stocks through inhibition of photosynthesis (Erftemeijer and Lewis, 2006) leading to major die back (Cyrus et al. 2008) and/or seagrass loss (Cabaço et al. 2008). The direct effects of extraction on seagrass meadows are when soil is removed to deepen and maintain navigation channels and harbour entrances or for extraction of fill materials from seagrass meadows (Larkum and West 1990; Da Silva et al. 2004). These activities have been reported to represent the greatest environmental threats to seagrass meadows, particularly in the Gulf region where more than 40% of the coastline has now been modified and developed (Sheppard and Price, 1991). The indirect effects of extraction can result from any significant dredging activity (i.e. beach renourishment, channel dredging or widening, changes in land catchments, etc) that occurs landward of seagrass meadows. The CO₂ emissions and removals associated with extraction (direct physical removal of biomass and soil) are similar to those of aquaculture during construction and so similar guidance is given).

Harvesting of aquatic resources

Mangrove Forests. Throughout the tropics, mangrove wood is harvested for fuel, charcoal, and construction (Ellison and Farnsworth 1996, Walters et al. 2008). Harvesting can range from extensive forest clearing, to more moderate, selective harvesting of individual trees to minimally invasive incidental bark removal. In wetlands dominated by woody vegetation change in biomass C stocks should be considered during both selective harvesting and forest clearance. CO₂ emissions from soils due to forest clearing should also be considered in C stock estimations (Alongi et al. 1998). To date, there have been no studies that suggest a change in N₂O emissions with harvests. Plant-mediated transport of methane has been documented (Kreuzwieser et al. 2003; Cheng et al. 2007), but few data are currently available.

Tidal Marshes. Harvesting of marsh vegetation is assumed to have negligible impacts on biomass and soil C stocks. However, in cases where extensive harvesting of vegetation or grazing is known to occur, it is good practice to document and report the CO₂ emissions and removals associated with this activity.

Seagrass Meadows. Physical disturbance is an issue with harvesting marine resources such as bivalves and other shellfish. On-ground shellfish culture and harvest methods can result in trampling and disturbance from boat wakes and propeller scars in shallow waters (Dumbauld et al. 2009). Intertidal seagrass beds can be particularly impacted by bait collection, through digging and trenching as well as pumping of sediment for prawns (Pillay et al. 2010) and associated trampling. Thus, harvesting of aquatic resources can have impacts on CO₂ emissions and removals primarily through changes in biomass C stocks where seagrass meadows have been lost.

Nutrient enrichment

Human activities can provide excess organic matter, nitrogen and phosphorus to coastal waters through agriculture runoff, discharge of sewage and industry effluent (Seitzinger and Harrison 2008). Nutrient

enrichment stimulates algal production, adding to organic matter loads in coastal waters. Nitrate, ammonium and dissolved organic N are the major forms of N input (Dai et al. 2008, Seitzinger and Harrison 2008). Decomposition of organic matter and nutrients supplied through anthropogenic sources can result in enhanced emissions of greenhouse gases.

Mangrove Forests and Tidal Marshes. In mangrove forests and tidal marshes, the effects nutrient availability on plant community structure and aboveground and belowground biomass is not well understood. Numerous fertilization experiments indicate that aboveground production in saline wetlands is limited by availability of N, while tidal freshwater marshes may be limited by availability of P (Megonigal and Neubauer 2009). Fertilization with N in tidal salt marshes has resulted in an increase in plant height, photosynthetic rates, and aboveground biomass production (e.g. Valiela and Teal 1974; Leendertse et al., 1997; Pennings et al., 2002). However, belowground production decreases with increased nutrient supply (Valiela et al., 1976; Darby and Turner 2008; Deegan et al., 2012). Since belowground production is the major source of carbon sequestered in marsh and mangrove soils, fertilization may result in no, or perhaps decreased, rates of carbon sequestration (Deegan et al. 2012). However, scientific evidence is not adequately developed to provide default emission factors for nutrient enrichment effects on CO₂ emissions and removals associated with biomass or soils in mangrove forests and tidal marshes. For instance, most mangrove research has focused on short-term increases in foliage nutrient content (e.g., Simpson et al., 2013) than on actual measurement of increases in biomass over time and the rate of increase in biomass increases initially with rate of nutrient supply (Alongi 2011). Thus, for the management activity nutrient enrichment, this chapter covers only CH₄ emissions due to effluent from aquaculture and urbanisation of the coastal zone and agricultural run-off and N₂O due to effluent from aquaculture. For non-CO₂ gases associated with other types of nutrient enrichment, covers N₂O emissions from agricultural run-off (through N leaching from application of N to terrestrial soils) are covered in Chapter 11, Volume 4 of the 2006 IPCC Guidelines. In this Supplement, N₂O and CH₄ emissions from waste water (including sewage) are covered in Chapter 6 on semi-natural treatment wetlands.

Seagrass Meadows. Decline of seagrass populations associated with anthropogenic nutrient loading including effluent from aquaculture (Yang and Yang, 2009) has been observed in many estuarine embayments (Waycott et al. 2009). Increased availability of nutrients may lead to blooms of macroalgae, phytoplankton, and epiphytes, all of which shade seagrass, reducing the light available for photosynthesis that can lead to seagrass loss (Waycott et al. 2009). Because the effects due to agricultural run-off and effluent discharge from aquaculture result in a decrease in both aboveground and belowground biomass stocks as well as increased CO₂ emissions from soils, with sufficient scientific evidence of both, CO₂ emissions and removals due to nutrient enrichment are covered in this Chapter.

Hydrologic/Sediment diversion (soil elevation loss)

Mangrove Forests and Tidal Marshes. The processes by which elevation in tidal wetlands is maintained against sea level rise and in some cases, coastal subsidence, rely upon the supply of mineral soils (Kirwan et al., 2011) derived from nearby rivers, cliffs, seabeds and oceanic import as well as growth of belowground biomass to maintain elevations against sea level rise. Therefore management practices that alter soil supply or cause subsidence will impact on coastal ecosystems. For example, many coastal wetlands are impacted by upstream diversions of water, such as from the construction of dams that is accompanied by diversion of sediment resulting in sediment starvation. Reduction of sediment delivery brought about by coastal structures can also result in sediment starvation, erosion of coastal wetland soils and release of stored carbon. Oil and gas extraction also affects coastal wetlands and results in soil subsidence and vegetation drowning in coastal areas (Craig et al. 1979; Dahl 2011). Thus hydrologic diversion from coastal areas with mangrove forests and tidal marshes, resulting in reduced elevation relative to sea-level rise or subsidence, can cause significant impacts to above and belowground C stocks if the soil becomes subtidal. As C in decayed vegetation is released to the water column and to the atmosphere, these C stock changes should be reported.

Drainage

Mangrove Forests and Tidal Marshes. Coastal wetlands have been diked and drained to create agricultural land (pasture and croplands) and settlement since before the 11th century (Gedan et al., 2009). The practice continues today on many coastlines. On some diked coasts, groundwater of reclaimed former wetlands is pumped out to maintain the water table at an optimum level below a dry soil surface while on other coasts drainage is achieved through a system of ditches and tidal gates. Drainage causes a reduction in the degree of soil saturation and ordinarily increases rates of organic matter decomposition, resulting in loss of soil carbon (Armentano and Menges 1986). This response will vary regionally with climate (Pozo and Colino 1992) and locally with soil salinity, soil texture, and the quantity of labile organic matter available in the drained soil (Heminga et al., 1998; Setia et al., 2011). Drainage can also lead to CO₂ emissions due to oxidation of dissolved organic and particulate

Second Order Draft

organic carbon in the water carried by drainage channels (see Annex 4.2). Drainage could be accompanied by land clearing, resulting in changes in biomass and soil carbon pools. If burning accompanies drainage, emissions from changes in those C pools (e.g. DOM) should also be reported.

Restoration and Creation.

Mangrove Forests and Tidal Marshes. Rewetting as a component of restoration, results where hydrologic modifications reverse drainage or remove impoundments or other obstructions to hydrologic flow (i.e. levee breach). Rewetting of drained soils through reconnection of hydrology can reduce oxidation of soil C but may lead to CH₄ emissions depending on salinity of reintroduced water (Harris et al 2010). The reestablishment of vegetation can reinitiate soil carbon sequestration leading to wetland restoration. Improved estimates of soil carbon accumulation rates in mangroves and tidal marshes (Chmura et al. 2003, Breithaup et al., 2012) have recently been obtained making it possible to quantify C gains in restored systems. Also included in this activity are created mangrove forests and tidal marshes that are typically established by removing the upper layer of upland soil or dredge spoil and grading the site until the appropriate tidal elevation is reached. Hence, the initial soils at created wetland sites are typically upland soils and/or dredge spoil. A peat layer can eventually develop in the surface soils with soil properties equivalent to soil properties within the upper layer of natural wetlands (Osland et al. 2012). Mangrove forests can also be created through high riverine sediment load, leading to rapid sediment accumulation, so that previously sub-aqueous soils can be elevated above tidal influence. This naturally created land can be naturally or purposefully vegetated. Time lags for equivalence of C pools between created and natural mangrove sites also suggest a timescale of ~20 years for restoration of functionally equivalent soil C accumulation rates (Craft et al. 2003, Osland et al. 2012).

Seagrass Meadows. Water quality improvements can reverse seagrass loss due to increased light penetration. The re-establishment of seagrass vegetation can increase C gains, albeit slowly. For example, a 3-5 year time lag between nutrient load reduction and initiation of seagrass recolonization has been observed (Vaudrey et al. 2010), with a further 12-15 yr time lag before seagrass biomass attains a relatively stable distribution with potential for soil carbon accumulation rates equivalent to natural settings (McGlathery et al., 2012). Restoration of sites previously supporting seagrass meadows can be mediated by purposeful dispersal of seed or planting of seagrass modules (Orth et al., 2012). These same techniques can also be used to create new seagrass meadows (Jones et al., 2012).

Other Activities.

Mangrove Forests, Tidal Marshes and Seagrass Meadows. In order not to omit any potentially significant management activity that may now or in the future constitute significant CO₂ emissions or removals or emissions of non-CO₂ gases, a final subtype “other activities” is identified. For example, currently salt production is currently not considered a management activity that significantly impacts seagrass meadows and thus no Tier 1 guidance is provided. However, as economic or social conditions change, so might management practices. Thus, the management activity termed, other activities, is provided to capture such potential changes in conditions.

4.1.2 Conversion from a land-use category that includes coastal wetlands

This section covers coastal wetland ecosystems that have been converted from the Wetlands land-use category to other land-use categories (see Section 4.3 for further guidance). Drainage or extraction are activities that are most likely to result in land-use category conversion from Wetlands to another IPCC land-use category, however, this is largely a function of how the land is classified (see Box 4.1). Excavation of soil may also result in conversion to Settlements. However, not all cases of land-use conversion result from drainage or extraction. For example, in the case of forested coastal wetland ecosystems, a land-use conversion can occur simply based on how the land is classified. For example, a mangrove forest that was cleared for timber harvest or otherwise deforested could undergo a change in land-use category if the mangrove forest had been classified as Forest land, but would not undergo a change in land-use category if the mangrove forest had been classified as Wetlands. Although the scale and resulting management activity may result in a land-use category change, the methodology to estimate CO₂ emissions is the same as given in section 4.1.1. There are cases in which a change in land-use category involves implementation of new methodologies and emission factors. For example, if a coastal wetland is drained to the extent that it is dry all year round, it does not meet the definition of wetlands as presented in this Chapter. Thus, a new land-use category would necessarily be applied. In the case that the coastal wetland has organic soils and is permanently drained wetland, refer to Chapter 2 of this Supplement (see

Box 4.1 for further examples). If conversion results in a land-use category with mineral soils, refer to the respective Chapter in the *2006 IPCC Guidelines*.

4.1.3 Conversion to a land-use category that includes coastal wetlands

This section considers rewetting and restoration or creation of coastal wetlands that occurs where hydrology, water quality or soils have been altered for establishment and survival of functioning vegetation (see Section 4.4 for further guidance). This includes restoration of coastal wetlands, where vegetation that was present prior to the current land use has been re-established, or creation, where actions have been taken to develop coastal wetlands in areas where they cannot be confirmed to have occurred previously.

4.2 LAND REMAINING IN A LAND-USE CATEGORY

This section details methodology necessary to estimate impacts of various human activities on greenhouse gas emissions from Land remaining in a land-use category. Sub-sections detail the methods for estimates as a result of change in biomass (subsection 4.2.1), DOM (subsection 4.2.2), soil (subsection 4.2.3), and for non-CO₂ gases (sub-section 4.2.4). Methods for biomass, DOM and non-CO₂ gases in large part reference the methodological guidance of the *2006 IPCC Guidelines* which should be read before and referred to in conjunction with, this guidance.

Guidance is provided here on estimating greenhouse gas emissions from management activities that do not result in a land-use conversion. These activities either take place without any changes of the water table over all land area (e.g. nutrient enrichment, harvesting of aquatic resources) or over part of the land area (e.g. salt production, extraction) within the managed coastal wetland. If these management activities can result in a change from the coastal wetland land-use category to another IPCC land-use category, then they will be considered in section 4.3.

The general approach that is applied combines the change in CO₂ emissions and removals for each activity summed for each pool with non-CO₂ gases treated separately. The following equation (Equation 4.1) is used for each of the carbon pools:

$$\begin{aligned} &\text{EQUATION 4.1} \\ &\text{ANNUAL CHANGE IN CARBON POOLS IN LAND REMAINING IN A LAND USE CATEGORY} \\ &\sum \Delta C_{\text{pool}(1...i)} = \\ &\Delta C_{\text{AQ}} + \Delta C_{\text{SP}} + \Delta C_{\text{EXT}} + \Delta C_{\text{HARV}} + \Delta C_{\text{NUTR}} + \Delta C_{\text{DIV}} + \Delta C_{\text{DR}} + \Delta C_{\text{RES}} + \Delta C_{\text{CRE}} \end{aligned}$$

where:

$\sum \Delta C_{\text{pool}(1...i)}$ = the sum of carbon stock changes for each pool with activities summed within each pool from 1 to i activities.

ΔC_{AQ} = Change in carbon stocks associated with aquaculture activity

ΔC_{SP} = Change in carbon stocks associated with salt production activity

ΔC_{EXT} = Change in carbon stocks associated with extraction activity

ΔC_{HARV} = Change in carbon stocks associated with harvesting activity

ΔC_{NUTR} = Change in carbon stocks associated with nutrient enrichment activity

ΔC_{DIV} = Change in carbon stocks associated with hydrologic/sediment diversion activity

ΔC_{DR} = Change in carbon stocks associated with drainage activity

ΔC_{RES} = Change in carbon stocks associated with restoration activity

ΔC_{CRE} = Change in carbon stocks associated with creation activity

4.2.1 Biomass

This section addresses estimation of changes in aboveground and belowground biomass carbon pools associated with management activities in coastal wetlands in all categories i.e. land remaining in a land-use category, conversion from a land-use category that includes coastal wetlands and conversion to a land-use category that includes coastal wetlands. It builds on the *2006 IPCC Guidelines* and the *Good Practice Guidance for Land Use, Land-Use Change and Forestry (GPG-LULUCF)*. For coastal wetlands with mangroves forest changes in biomass carbon stocks follow equations provided in Chapters 2 and 4, Volume 4, of the *2006 IPCC Guidelines*. For managed coastal wetlands with tidal marshes and seagrass meadows, changes in biomass carbon stocks follow largely follow equations in the Chapters 2 and 6, Volume 4, of the *2006 IPCC Guidelines*. However, there are some deviations from these general methods as applied to the Land Remaining in a land-use category guidance as presented in Volume 4 of the *2006 IPCC Guidelines*. Thus, Table 4.2 should be used to identify the appropriate Tier 1 approaches and equations that should be applied when following this guidance. The main Tier 1 approaches and equations for each activity are presented in Table 4.2 below.

TABLE 4.2 MANAGEMENT ACTIVITIES, TIER 1 EQUATIONS OR DEFAULT EF FOR BIOMASS POOL CHANGES		
Activity	Ecosystem	Tier 1 Equation ¹ and Default EF
Aquaculture (ΔC_{B-AQ}), Salt Production (ΔC_{B-SP}), Extraction (ΔC_{B-EXT})	Mangrove forest	AQ, SP, EXT-D² : at start of activity, apply Eq. 2.9 or 2.10 and Eq. 2.11 & 2.14; $\Delta C_G = 0$ and $\Delta C_L = L_{\text{disturbance}}$; Table 4.3 – 4.6 AQ, SP : for use phase, $\Delta C_G = \Delta C_L$; Table 4.3-4.6 AQ, SP : for abandonment phase, apply Eq. 2.9 or 2.10 and Eq. 2.11 - 2.14; Table 4.3-4.7
	Tidal marsh	AQ, SP, EXT-D² : at start of activity, apply Eq. 2.8a where $C_{t2}=0$ and C_{t1} =initial C stock ³ ; Table 4.8 AQ, SP : for abandonment phase, apply Eq. 2.8a; Table 4.8 AQ : for use phase, $\Delta C_G = \Delta C_L$; Table 4.8
	Seagrass meadow	AQ, EXT-D and EXT-I⁴ : at start of activity, apply Eq. 2.8a where $C_{t2}=0$ and C_{t1} =initial C stock ³ ; Table 4.9-4.11
Harvesting of Aquatic Resources (ΔC_{B-HARV}), Hydrologic/Sediment Diversion (ΔC_{B-DIV})	Mangrove	Apply Eq. 2.9 or 2.10 and Eq. 2.11 - 2.14 ⁵ ; Table 4.3-4.6
	Tidal Marsh	Apply Eq. 2.8a; Table 4.8
	Seagrass Meadow	NA
Nutrient Enrichment (ΔC_{B-NUTR})	Mangrove	NA
	Tidal Marsh	NA
	Seagrass Meadow	EXT-I⁴ : at start of activity, apply Eq. 2.8a where $C_{t2}=0$ and C_{t1} =initial C stock ³ ; Table 4.9-4.11
Drainage (ΔC_{B-DR}), Restoration (ΔC_{B-RES}) & Creation (ΔC_{B-CRE}), Other Activities ($\Delta C_{B-OTHER}$)	Mangrove	DR, RES, CR, OTHER : apply Eq. 2.9 or 2.10 and Eq. 2.11 - 2.14; Table 4.3 – 4.7
	Tidal Marsh	DR, RES, CR, OTHER : apply Eq. 2.8a; Table 4.8
	Seagrass Meadow	DR: NA, RES, CR, OTHER : apply Eq. 2.8a; Table 4.9-4.11
Note -- ¹ Equations in the section can be found in the 2006 IPCC Guidelines; ² EXT-D = where biomass are physically removed; ³ C_{t1} =initial C stock can be found in Table 4.9-4.11; ⁴ EXT-I = biomass C stock change resulting from sediment deposition effects; ⁵ reporting $L_{\text{disturbance}}$ as loss due to harvest; NA = not applicable		

4.2.1.1 CHOICE OF METHOD

Removal or loss of biomass resulting from management activities that occur during aquaculture, salt production, and extraction for mangrove forests, tidal marshes and seagrass meadows are estimated at Tier 1 level with the

key assumption that carbon in biomass lost in the year of the start of the activity is oxidised to CO₂ (Table 4.2). For mangrove forests, this is estimated with the *Gain-Loss* method as $\Delta C_L = L_{\text{disturbance}}$. For tidal marshes and seagrass meadows, this is estimated with the *Stock-Difference* method with the biomass C stock at $t_2 = 0$. For harvesting of aquatic resources and hydrologic/sediment diversion, guidance follows equations provided in Chapter 2 as noted in Table 4.2. For nutrient enrichment, change in biomass C stocks are only estimated for seagrass meadows as EXT-I (indirect effect of extraction activity; see Section 4.1.1) and follows the *Stock-Difference* approach with the biomass C stock at $t_2 = 0$ (lost in the year that the activity starts). For drainage, restoration/creation and other activities, methodology follows guidance in *2006 IPCC Guidelines* as described in Table 4.2 with the exception of drainage for seagrass meadows, an activity for which there is no evidence of its occurrence.

TIER 1

Mangrove, Tidal Marsh and Seagrass Meadow

The method for estimating changes in mangrove biomass for Tier 1 follows Eq. 2.9 and Eq. 2.11-2.14 and for tidal marsh and seagrass biomass Eq. 2.8a is applied (see Volume 4 of the *2006 IPCC Guidelines* for details). Default values presented in Tables 4.3-4.11 are used in these equations, depending on whether the ecosystem is mangrove forest, tidal marsh, or seagrass meadow and their climate domain. For mangroves, Tables 4.3-4.7 contain default factors for average aboveground biomass, the ratio of belowground to aboveground biomass, and dry weight:carbon conversion factors, differentiated by humid and dry tropical regions and the subtropics. To estimate losses due to wood-removals and fuelwood, wood density (Table 4.7) and BEF values should be applied (Chapter 3 in 2003 *GPG-LULUCF*). For tidal marshes, Table 4.8 contains default factors for average aboveground and belowground biomass by climate and salinity. For seagrass meadows, methodology also generally follows the *Stock-Difference* approach with data provided for aboveground biomass, the ratio of belowground to aboveground biomass and C conversion factors (Tables 4.9-4.11). It should be noted that the loss of biomass C stock from seagrass meadows due to nutrient enrichment and indirect effects of extraction results in loss of water clarity and light available (as described in Section 4.1.1) and should be reported.

TIER 2

Tier 2 methods are used where countries have country-specific emission factors, and substantial national data or data that can be gathered based at reasonable cost. Country-defined methodology (e.g., high priority is empirical measurement of above-ground biomass which for mangroves involves use of allometric equations) may be based on detailed inventories of permanent sample plots for coastal wetlands. Tier 2 can include such other improvements as, in the case of mangroves, use of allometric relationships to accurately measure aboveground biomass; a priority in this case would be to measure diameter-at-breast height of trees in the permanent plots (see Annex 4.3). To estimate carbon gain using country-specific annual increment, wood density (Table 4.7) and BEF values should be applied (Chapter 3 in 2003 *GPG-LULUCF*). In the case of seagrasses, where extraction, nutrient enrichment, or aquaculture resulting in either reduced water quality or redeposition of sediments is significant leading to a land category or pool that is key, Tier 2 methods can involve data collected from permanent sample plots or other means of acquiring country-specific data as covered in standard texts. For seagrass meadows, the lag time between water clarity impacts and biomass C stock change could be employed where t_2 was estimated as a fraction of initial stock.

TIER 3

Tier 3 approach for biomass carbon stock change estimation allows for a variety of methods, including process based models. Implementation may differ from one country to another, due to differences in inventory methods, coastal wetland conditions and activity data. Tier 3 requires use of detailed national coastal wetland inventories. They may be supplemented by allometric equations and models calibrated to national circumstances that allow for direct estimation of biomass growth. Tier 3 could also involve inventory systems using statistically-based sampling of biomass over time and/or process models, to further stratify ecosystem type into subtype, ecological zone and salinity.

4.2.1.2 CHOICE OF EMISSION/REMOVAL FACTORS

TIER 1

Mangrove, Tidal Marsh and Seagrass Meadow

Second Order Draft

Countries using a Tier 1 method should utilize default data for mangrove forests, tidal marshes or seagrass meadows provided in this guidance. For mangrove forests, prior to when or where aquaculture, salt production, extraction or harvesting, or hydrologic/sediment diversion occurred, Tier 1 emission factors for biomass stocks can be calculated using a *Stock-Difference* approach or a *Gain-Loss* approach using biomass growth, primarily differentiated at the level of climate domain (and in the case of tidal marshes, also by salinity for the temperate climate domain). For mangrove forests, the use of Tables 4.3 – 4.7 including guidance on estimating BCEF based on woody density for losses of fuelwood and wood-removals found in the *2006 IPCC Guidelines* are needed. For tidal marshes, the default emissions factors to be used are in Tables 4.8. For seagrass, emission factors found in Tables 4.9-4.11.

TIER 2

The higher Tier methods described in Chapter 2 of the *2006 IPCC Guidelines* (Eq. 2.18-2.23) will permit better estimates when national data are applied. Estimate of loss of below-ground root biomass should also be empirically measured rather than using the ratio of aboveground to belowground biomass.

TIER 3

Tier 3 modelling would include country-specific emission factors derived or modelled from measurement data. The models should capture variation in emission rates driven by extent and depth of biomass extraction and in the case of water/sediment diversion, the rate of biomass decline. The incremental growth rates of the dominant species can be multiplied by the current biomass estimate to derive a figure of projected change over time, if required. For seagrass meadows, aerial photographs may be used in regions of high water clarity, good contrast between seagrass meadows and soil and low cover of other macrophytes. Field validations can be implemented to verify model output using field measurements by diving or underwater camera. Other examples of data improvements for aquaculture, salt production and extraction, for example, include ground-truth estimates of total area impacted, the depth at which removal of biomass has occurred or the completeness of biomass removal could also be verified. Furthermore, with indirect effects of extraction on seagrass meadows, because the time periods for biomass loss will vary based on national circumstances, some seagrass meadows will show more rapid decline than seagrass meadows in other areas. Total areas can be estimated with greatest certainty coupling aerial or remote sensing with field measurements.

4.2.1.3 CHOICE OF ACTIVITY DATA

All tiers require information on areas of managed coastal wetlands for ecosystem types, climate, and management activities which vary depending on national circumstances. The default methodology for all Tier 1 estimates requires data on the area of managed wetland, the type and dominant species present in the wetland, and soil type. The type of wetland present as well as soil type can be obtained from national wetland and soil type maps (if available) or the International Soil Reference and Information Centre; www.isrig.org; wetland distributions for most countries can be obtained from the RAMSAR web site (www.ramsar.org). When information is gathered from multiple sources, cross-checks should be made to ensure complete and consistent representation and avoid omissions and double-counting. Please refer to the extended list of activity data sources in Annex 4.2 in this Chapter.

Aquaculture, Salt production

The ponds or structures used to create conditions conducive to aquaculture and salt production can be identified from aerial photographs or by inspection and measurement of the total area where biomass has been removed. Information on regional aquaculture projects worldwide can be obtained from the GISFish database of the FAO (www.fao.org) and similar project information for salt production projects can be obtained from the Salt Institute at www.saltinstitute.org.

Extraction or Harvesting

Information of current extraction/harvesting projects may be obtained from local, state or regional government department websites as many countries and regional government authorities require documentation of environmental impact assessments. For seagrass meadows, aerial photographs may be used in regions of high water clarity, good contrast between seagrass and soil and low cover of other macrophytes. If these conditions are not present, remote sensing imagery providing data on water clarity using intensity of chlorophyll or concentration of suspended particles other photoimagery can be used and readily available.

Nutrient enrichment

We can expect to find nutrient-affected systems downstream of major urban centers, municipal sewage treatment facilities, and watersheds where a large portion of the land is under intensive agricultural use. For seagrass meadows the N load per estuary that relates to complete loss of seagrass has been predicted to be at loading values between 100 and 175 kg N ha⁻¹ yr⁻¹ (Steward and Green 2007; Latimer and Rego 2010). Information about changes in seagrass biomass as a result of nutrient enrichment may be obtained from national government departments, marine conservation trusts and societies and global seagrass networks (<http://seagrasswatch.org>). Review papers in the scientific literature (e.g. Waycott et al., 2009) have collated information on localised seagrass loss. Historical records of seagrass cover including aerial photographs may be used in regions of high water clarity, good contrast between seagrass and soil and low cover of other macrophytes.

Hydrologic/Sediment Diversion, Drainage

Total areas impacted by upstream diversions of water and sediment result in soil elevation loss and coastal subsidence can be assessed using aerial or remotely sensed data as well as reports from Environmental ministries or other agencies reporting rates of coastal subsidence and at higher Tiers validated with field measurements. Drainage is recognized by presence of levees and rectilinear channel networks that stand out in comparison to the normally flat surface with curving, bifurcating channels. Restoration would be recognized by the removal of levees, tidal gates, enlargement of culverts, and possibly development of new hybrid channel networks that incorporate both constructed and natural channels (MacDonald et al., 2010). Channels may visibly widen. Immediately after rewetting, dead vegetation may be visible in tidal marshes, in both mangroves and tidal marshes fresh mud will be deposited over previous soil surface. These information can be easily detected on aerial images that are freely available. If following Approach 1 for land classification, information of drainage and diversion projects may be obtained from local, state or regional government department websites as many countries and regional government authorities require documentation of environmental impact assessments for these activities.

Restoration/Creation

Estimates of biomass change as a result of wetland restoration or creation requires knowledge of previous wetland coverage (in the case of restoration) and of biomass growth rates (in the case of restoration or creation). Historical photos may be used to most readily estimate the pre-restored wetland area. Information on regional wetland restoration and creation projects worldwide can be obtained from the Global Gateway to Geographic Information Systems of the FAO (www.fao.org) as well as from the websites, www.wetlands.org and www.globalrestorationnetwork.org.

Table 4.3 Carbon content of aboveground mangrove forest biomass ((gC/100gDW) or % DW)

Component	%C	95% CI	Range	References
Leaves	45.0 (n = 35)	42.8, 47.3	42.3-50.9	Spain and Holt, 1980; Saenger, 2002; Alongi et al., 2003; 2004; Kristensen et al., 2008
Wood (stems + branches)	45.4 (n = 12)	43.1, 47.7	41.6-47.2	Spain and Holt, 1980; Gong and Ong, 1990; Twilley et al., 1992; Bouillon et al., 2007; Alongi et al., 2003, 2004

Table 4.4 Aboveground Biomass in Mangrove forests (tonnes DW. ha⁻¹)

Domain	Region	Aboveground biomass	95%CI	Range
Tropical	Tropical Wet	196 (n=53) ¹	186, 205	3.7-557
	Tropical Dry	92 (n = 13) ²	88, 97	3.2-201
Subtropical		75 (n= 10) ³	76, 84	3.9-129

¹References: Golley et al., 1975; Christensen, 1978; Ong et al., 1982; Putz and Chan, 1986; Tamai et al., 1986; Komiyama et al., 1987, 1988, 2000, 2008; Lin et al., 1990; Mall et al., 1991; Amarasinghe and Balasubramaniam, 1992; Kusmana et al., 1992; Slim et al., 1996; Fromard et al., 1998; Norhayati and Latiff, 2001; Pongparn, 2003; Sherman et al., 2003; Juliana and Nizam, 2004; Kirui et al., 2006; Kairo et al., 2008; Fatoyinbo et al. 2008; Camacho et al., 2011; Kauffman et al., 2011; Thant and Kanzaki, 2011.

²References: Golley et al., 1962; Briggs, 1977; Suzuki and Tagawa, 1983; Steinke et al., 1995; Alongi et al., 2003; Medeiros and Sampoia, 2008; Khan et al., 2009.

³References: Lugo and Snedaker, 1974; Woofroffe, 1985; Lee, 1990; Mackey, 1993; Tam et al., 1995; Saintilan, 1997; Ross et al., 2001; Coronado-Molina et al., 2004; Simard et al., 2006; Fatoyinbo et al., 2008; Komiyama et al., 2008; Abohassan et al., 2012.

Second Order Draft

Table 4.5 Aboveground biomass growth in mangrove forests (tonnes DW ha⁻¹ yr⁻¹)

Domain	Region	Aboveground biomass growth	95%CI	Range
Tropical	Tropical Wet	9.9 (n=23) ¹	9.4, 10.4	0.1-27.4
	Tropical Dry	3.3 (n = 6) ¹	3.1, 3.5	0.1-7.5
Subtropical		18.1 (n= 4) ¹	17.1, 19.1	5.3-29.1

¹References: Ajonina 2008; Kairo et al., 2008; Alongi 2010

Table 4.6 Ratio of belowground biomass to aboveground biomass (R) in mangroves forests

Domain	Region	R	95%CI	Range
Tropical	Tropical Humid	0.49 (n=18) ¹	0.47, 0.51	0.04-1.1
	Tropical Dry	0.29 (n = 9) ²	0.28, 0.30	0.09-0.79
Subtropical		0.96 (n= 18) ³	0.91, 1.0	0.22-267

¹References: Golley et al., 1975; Tamai et al., 1986; Komiyama et al., 1987, 1988; Gong and Ong, 1990; Lin and Lu 1990; Tam et al., 1995; Pongpam, 2003

²References: Golley et al, 1962; Alongi et al., 2003; Hoque et al., 2010.

³References: Briggs, 1977; Lin, 1989; Saintilan, 1997.

TABLE 4.7. WOOD DENSITY (D) OF COMMON MANGROVE TREE SPECIES

<i>Bruguiera gymnorrhiza</i>	0.74	<i>Heritiera littoralis</i>	0.8
<i>Xylocarpus mekongensis</i>	0.72	<i>Heritiera fomes</i>	1.1
<i>Xylocarpus granatum</i>	0.7	<i>Excoecaria agallocha</i>	0.45
<i>Sonneratia apetala</i>	0.56	<i>Ceriops tagal</i>	1.1
<i>Sonneratia alba</i>	0.08	<i>Ceriops decandra</i>	0.96
<i>Rhizophora mucronata</i>	1.1	<i>Bruguiera gymnorrhiza</i>	0.86
<i>Rhizophora mangle</i>	0.83	<i>Avicennia officinalis</i>	0.67
<i>Rhizophora apiculata</i>	1.1	<i>Avicennia marina</i>	0.9
<i>Laguncularia racemosa</i>	0.6	<i>Avicennia germinans</i>	0.66
Site average of above values	0.77		
Data sources: Kauffman and Donato, 2012; Bosire et al., 2012.			

651

TABLE 4.8 ABOVEGROUND AND BELOWGROUND BIOMASS FOR TIDAL MARSHES (TONNES C HA ⁻¹)								
Salinity	Domain	Region	Above-ground biomass	95% CI	n	Below-ground biomass	95% CI	n
Oligohaline to polyhaline	Subtropical	Warm temperate moist ¹	1.40	1.35, 1.44	3	9.2	8.58, 9.80	2
		Warm temperate dry ²	5.64	5.57, 5.71	11	11.02	10.89, 11.15	4
	Temperate ³		3.32	3.30, 3.34	91	23.9	23.37, 24.32	34
	Polar ⁴		0.64	0.62, 0.66	3	ND ⁶		
Tidal freshwater	Temperate ⁵		4.03	4.00, 4.07	20	7.11	6.94, 7.18	8
Sources: ¹ AG: da Cunha Lana et al. 1991; Kirby and Gosselink 1976; Darby and Turner 2008; BG: da Cunha Lana et al. 1991; Darby and Turner 2008 ² AG: Curreo et al. 2002; Neves et al. 2007; Scarton et al. 2002; Linthust and Reimold 1978; Cartaxana and Catarina 1997; BG: Neves et al. 2007; Scarton et al. 2002; also referred to as Mediterranean ³ AG: Benito and Onaindia 1991; Bouchard and Lefevre 2000; Boyer et al. 2000; Connor and Chmura 2000; Figueroa et al. 1988; Gallagher and Plumley 1979; Groenendijk 1984; Gross et al. 1986; Hopkinson et al. 1978; Hussey and Long 1982; Jeffrey and Hayes 2005; Jeffries et al. 1981; Jenssen 1980; Keefe and Boynton 1973; Ketner 1972; Kirby and Gosselink 1976; Kistritz et al. 1983; Linthust and Reimold 1978; Mahall and Park 1976; Mendelsohn and Marcellus 1976; Morris and Haskin 1990; Pierce 1983; Schubauer and Hopkinson 1984; Smith et al. 1979; Squires and Good 1974; Whigham et al. 1978; White et al. 1978; Williams and Murdoch 1972; BG: Boyer et al. 2000; Connor and Chmura 2000; Dame and Kenny 1986; Dunn 1981; Elsey-Quirk et al. 2011; Groenendijk and Vink-Lievaart 1987; Gross et al. 1991; Hussey and Long 1982; Kistritz et al. 1983; Livingstone and Patriquin 1981; Mahall and Park 1976; Roman and Daiber 1984; Smith et al. 1979; Whigham et al. 1978 ⁴ AG: Curreo et al. 2002; Neves et al. 2007; Scarton et al. 2002; Linthust and Reimold 1978; Cartaxana and Catarina 1997; BG: Neves et al. 2007; Scarton et al. 2002 ⁵ Birch and Cooley 1982; Doumlele 1981; Hopkinson et al. 1978; Whigham et al. 1978 ⁶ No data available								

652

653

654

655

656

657

658

659

660

661

662

663

664

665

666

Second Order Draft

TABLE 4.9 ABOVEGROUND BIOMASS FOR SEAGRASS MEADOW

Domain	Aboveground biomass for seagrass meadow (tonnes DW ha ⁻¹)	95%CI	Range	n
Tropical ¹	6.0	6.20, 7.89	0.1 – 75.4	137
Subtropical ²	15.2	14.17, 16.26	1.2 – 810	260
Temperate ³	19.9	18.16, 21.77	4.3 – 206	108

¹Aioi & Pollard 1993, Brouns 1985, Brouns 1987b, Brouns & Heijs 1986, Daby 2003, de Jongh et al. 1995, Devereux et al. 2011, Fourqurean et al. 2012, Heijs 1984, Hertbert 1986, Holmer et al. 2001, Ismail 1993, Kaldy & Dunton 2000, Lee 1997, Lindeboom & Sandee 1989, Long et al. 1993, McDermid & Edward 1999, McKenzie 1994, Mellors et al. 2002, Moriarty et al. 1990, Nienhuis et al. 1989, Ogden & Ogden 1982, Paynter et al. 2001, Poovachiranon & Chansang 1994, Povidisa & Delefosse 2009, Rasheed 1999, Terrados et al. 1998, Uku & Bjork 2005, van Lent et al. 1991, van Tussenbroek 1998, Vermaat et al. 1993, Vermaat et al. 1995, Williams 1987.

²Aioi 1980, Aioi et al. 1981, Asmus et al. 2000, Bandeira 1997, Bandeira 2002, Boon 1986, Bulthuis & Woelkerling 1983, Burd & Dunton 2001, Cabaco et al. 2012, Calleja et al. 2005, Cambridge & Hocking 1997, Carruthers 1994, Collier et al. 2009, de Boer 2000, Dillon 1971, Dixon & Leverone 1995, Dos Santos et al. 2012, Dunton 1990, Dunton 1994, Dunton 1996, Fourqurean et al. 2012, Goshima & Peterson 2012, Heck & Thoman 1984, Heijs 1984, Herbert & Fourqurean 2009, Herbert & Fourqurean 2008, Hillman et al. 1995, Holmer & Kendrick 2012, Holmer et al. 2004, Iverson & Bittaker 1986, James et al. 2009, Kaldy & Dunton 2000, Kenworthy & Thayer 1984, Kerr & Strother 1989, Kim et al. 2012, Kirkman & Cook 1987, Kirkman & Reid 1980, Kirkman & Reid 1979, Kirkman et al. 1982, Kowalski et al. 2009, Kraemer & Alberte 1993, Larkum et al. 1984, Lee & Dunton 1996, Lee et al. 2005, Lewis 1987, Longstaff et al. 2000, Marba & Walker 1999, Masini et al. 2001, McGlathery et al. 2012, McMahon 1968, McMahon & Lavery 2008, Meling-Lopez & Ibarra-Obando 1999, Moncreiff et al. 1992, Morgan & Kitting 1984, Mukai et al. 1979, Murray and Wetzel 1987, Nienhuis & Bree 1980, Odum 1963, Orth & Moore 1986, Paling & McComb 2000, Park et al. 2011, Perez-Llorens & Niell 1993, Perez-Ruzafa et al. 2012, Phillips et al. 1983, Powell 1989, Preen 1995, Reyes et al. 1995, Schwarz et al. 2006, Stevensen 1988, Terrados & Ros 1992, Thayer et al. 1977, Tomasko & Lapointe 1991, Townsend & Fonseca 1998, van Houte-Howes et al. 2004, van Lent et al. 1991, van Tussenbroek 1998, Walker 1985, Walker & McComb 1988, Walker et al. 1988, West & Larkum 1979, Yarbro & Carlson 2008, Zavodnik et al. 1998, Zieman et al. 1989.

³Auby & Labourg 1996, Bay 1984, Cebrian et al. 1997, Cebrián et al. 2000, Dennison & Alberte 1985, Duarte et al. 2002, Fourqurean et al. 2012, Guidetti et al. 2002, Gullstrom et al. 2012, Hebert et al. 2007, Jacobs 1979, Kaldy 2006, Kentula & McIntire 1986, Larned 2003, Laugier et al. 1999, Lillebo et al. 2006, Mann 1972, Marba & Duarte 2001, Martin et al. 2005, McRoy 1974, Nienhuis & Bree 1980, Nixon & Oviatt 1972, Olesen 1999, Olesen & Sand-Jensen 1993, Olesen & Sand-Jensen 1994, Ott 2008, Pedersen & Borum 1993, Perez & Romero 1994, Plus et al. 2003, Rismondo et al. 1997, Robertson & Mann 1984, Robertson & Mann 1974, Roman & Able 1988, Sand-Jensen & Borum 1983, Sfriso & Ghetti 1998, Terrados et al. 2006, Vermaat & Verhagen 1996, Vermaat et al. 1987, Zavodnik et al. 1998.

TABLE 4.10 RATIO OF BELOWGROUND BIOMASS TO ABOVEGROUND BIOMASS (R) FOR SEAGRASS MEADOW

Domain	R	95%CI	Range	n
Tropical ¹	1.7 ¹	1.62, 1.78	0.05 – 25.62	396
Subtropical ²	2.4 ²	2.33, 2.54	0.07 – 16.8	391
Temperate ³	1.3 ³	1.18, 1.39	0.14 – 13.8	91

¹Aioi & Pollard 1993, Brouns 1985, Brouns 1987, Coles et al. 1993, Daby 2003, Devereux et al. 2011, Fourqurean et al. 2012, Halun et al. 2002, Holmer et al. 2001, Ismail 1993, Lee 1997, Lindeboom & Sandee 1989, McKenzie 1994, Mellors et al. 2002, Moriarty et al. 1990, Nienhuis et al. 1989, Ogden & Ogden 1982, Paynter et al. 2001, Poovachiranon & Chansang 1994, Povidisa et al. 2009, Rasheed 1999, Udy et al. 1999, van Lent et al. 1991, van Tussenbroek 1998, Vermaat et al. 1993, Vermaat et al. 1995, Williams 1987.

²Aioi 1980, Aioi et al. 1981, Asmus et al. 2000, Bandeira 2002, Boon 1986, Brun et al. 2009, Collier et al. 2009, de Boer 2000, Devereux et al. 2011, Dixon & Leverone 1995, Dos Santos et al. 2012, Dunton 1996, Fourqurean et al. 2012, Hackney 2003, Herbert and Fourqurean 2009, Herbert & Fourqurean 2008, Holmer & Kendrick 2012, Jensen & Bell 2001, Kim et al. 2012, Kirkman & Reid 1979, Kowalski et al. 2009, Larkum et al. 1984, Lee et al. 2005, Lee et al. 2005b, Lipkin 1979, Longstaff et al. 1999, Masini et al. 2001, McGlathery et al. 2012, McMahon 1968, Meling-Lopez & Ibarra-Obando 1999, Mukai et al. 1979, Paling & McComb 2000, Park et al. 2011, Powell 1989, Preen 1995, Schwarz et al. 2006, Stevensen 1988, Townsend & Fonseca 1998, Udy & Dennison 1997, van Houte-Howes et al. 2004, van Lent et al. 1991, van Tussenbroek 1998, Walker 1985, West & Larkum 1979, Yarbro & Carlson 2008.

³Agostini et al. 2003, Cebrian et al. 2000, Fourqurean et al. 2012, Hebert et al. 2007, Holmer & Kendrick 2012, Larned 2003, Lebreton et al. 2009, Lillebo et al. 2006, Marba & Duarte 2001, McRoy 1974, Olesen & Sand-Jensen 1994, Rismondo et al. 1997, Sand-Jensen & Borum 1983, Terrados et al. 2006

Table 4.11 Carbon content of aboveground and belowground seagrass biomass ((gC/100gDW) or % DW)				
Component	CF	95% CI	Range	References
Above ground biomass	33.4 (n = 72)	32.9, 33.9	23.6 – 43.8	Augier et al. 1982; Erftemeijer 1994; Fourqurean & Zieman 1992; Fourqurean et al. 1997; Fourqurean et al. 2007; Kenworthy & Thayer 1984; Longstaff & Dennison 1999; Mascaro et al. 2009; Mateo & Romero 1997; Nienhuis et al. 1989; Pergent et al. 1994; Plus et al. 2001; Rublee & Roman 1982; van Lent et al. 1991; Vinther & Holmer 2008; Wahbeh 1988.
Below ground biomass	32.3 (n = 50)	31.8, 32.9	24.7 – 42.1	Erftemeijer 1994; Kenworthy & Thayer 1984; Mascaro et al. 2009; Mateo & Romero 1997; Nienhuis et al. 1989; Plus et al. 2001; van Lent et al. 1991; Vinther & Holmer 2008; Wahbeh 1988.

4.2.1.5 UNCERTAINTY ASSESSMENT

This section considers source-specific uncertainties relevant to inventory estimates. Estimating country-specific and/or disaggregated values requires more accurate information on uncertainties than given below. Chapter 3, Volume 1 of the *2006 IPCC Guidelines* provides information on uncertainties associated with sample-based studies. The literature available on uncertainty estimates on emission factors and activity data is limited. Seagrass aboveground biomass averages values from species that have very different leaf characteristics, resulting in a wide range of values. In the case of mangrove aboveground biomass, a comparatively large number of forest sites are extant, but the range of values is very large due to inherent differences in species composition, forest age, tidal height, and soil fertility. These uncertainties may be minimized as much as possible using Tier 2 or Tier 3 methods which can incorporate or classify forests by factors other than climate domain.

Variability in tidal marsh biomass will be due to differences in dominant species and competition between species, as well as salinity of flood waters, frequency of tidal flooding and climate. The high biomass in Mediterranean climates is due to the frequent dominance of perennial shrubs. For all vegetation there can be considerable yearly variability in production of biomass and seasonal variability in standing biomass. The empirical data available is biased by the prevalence of data from temperate regions and North America - the tidal fresh marsh data comes almost entirely from the Atlantic coast of the United States. Also lacking are data from boreal and subtropical marshes.

EMISSION AND REMOVAL FACTORS

The major sources of uncertainty for all wetland types are dominant species-specific differences in carbon content and differences as a function of ecosystem age, species composition, intertidal location, and community structure. As indicated in Tables 4.3 to 4.11, coefficients of variation range from about 24-200%. To reduce uncertainty, countries are encouraged to develop country- or region specific biomass expansion factors and BCEFs that fit their conditions. In case country- or regional-specific values are unavailable, the sources of default parameters should be checked and their correspondence with species present and specific conditions of a country should be examined.

The causes of variation of annual increment of mangrove growth include climate, site growth conditions, and soil fertility. Artificially regenerated and managed stands are less variable than natural forests. The major ways to improve accuracy of estimates of these wetlands are by application of country-specific or regional estimates of growth stratified by the dominant species present. If the default values of growth increments are used, the uncertainty of estimates should be clearly indicated and documented.

Tier 3 approaches can use growth curves stratified by species, ecological zones, site productivity and management intensity. For example, similar approaches are routinely used for mangroves in timber supply planning models and this information can be incorporated into carbon accounting models (e.g., Kurz et al., 2002).

For mangroves, data on commercial fellings are relatively accurate, although they may be incomplete or biased due to illegal fellings and underreporting due to tax regulations. Traditional wood that is gathered and used directly, without being sold, is not likely to be included in any statistics. Countries must carefully consider these

Second Order Draft

issues. The amount of wood removed from forests after storm breaks and pest outbreaks varies both in time and volume. No default data can be provided on these types of losses. The uncertainties associated with these losses can be estimated from the amount of damaged wood directly withdrawn from the forest or using data on damaged wood subsequently used for commercial and other purposes. If fuelwood gathering is treated separately from fellings, the relevant uncertainties might be high, due to high uncertainty associated with traditional gathering.

4.2.2 Dead Organic Matter

The dead organic matter (DOM) pool in coastal wetlands includes coarse woody debris, fine litter, and dead roots. Fine litter and dead wood are differentiated as fractions <10 cm and >10 cm diameter, respectively (Table 1.1, Chapter 1, Volume 4 of the *2006 IPCC Guidelines*). The dead roots less than 2 cm diameter are included in the soil pool and not considered within the DOM pool. This fraction of dead roots turns over rapidly with the assumption approximating steady state. DOM C stocks can vary depending on tidal inundation and frequency, as well as soil oxidation and vegetation cover. Fine litter can be exported with tidal activity while a larger fraction of senesced woody biomass is buried or decomposes *in-situ*. In wetlands, decomposition of DOM, especially wood and coarse woody debris, is slow and accumulates as soil organic matter. In estimating inputs, outputs or changes of DOM C stocks, careful consideration of pools should be made to avoid double-counting. Consistency in how these fractions are determined and reported is good practice to avoid double-counting. DOM pools under conditions of low soil oxidation-reduction potential that occur under saturated conditions can be large and when exposed to oxidation constitute large CO₂ emissions. Thus, DOM C stock changes should be considered for the activities as described above with changes in mangrove forest cover. This is a notable deviation from Tier 1 assumptions for Land Remaining in a Land-Use Category as laid out in the *2006 IPCC Guidelines* and is necessary as certain types of management activities have large impacts on CO₂ emissions and removals without resulting in a conversion to another IPCC land-use category.

General assumptions for estimating changes in the DOM pool can be differentiated between mangrove forest and primarily non-woody coastal wetland types including tidal marsh and seagrass meadow. At Tier 1, tidal marsh and seagrass meadow ecosystems are assumed to have relatively fast turnover of DOM and a steady-state assumption applies. However, extensive management activities that result in vegetation or soil disturbance in tidal marsh with perennial biomass could have large impacts on C emissions and removals. In Land remaining in a land-use category, management activities where changes in the DOM pool are likely to be significant include aquaculture, salt production, extraction, where the start of the activity results in loss of the DOM pool. Methodologies for mangrove forests are supplied for Tier 1 estimation for management activities include aquaculture, salt production, and extraction (Table 4.1). In these cases, DOM inputs may cease as extant vegetation is cleared and DOM outputs are the extant stock of DOM. For drainage, DOM inputs may be unchanged if there is no disturbance to the biomass pool and DOM outputs are equivalent to the stock of DOM at the start of the activity where all C is assumed oxidized at Tier 1. For the harvesting activity and all other activities that affect mangrove forests, generic equations provided in *2006 IPCC Guidelines* are directly applicable. For all management activities that occur in tidal marsh and seagrass meadow, the change in C stocks of DOM is assumed to be zero at Tier 1 (Table 4.12). If mangrove forest clearing is a significant activity, Tier 2 or 3 approaches should be used and the assumption of instantaneous oxidation of the DOM pool is not valid. Both changes in dead wood and litter should be reported and summed to obtain changes in total dead organic matter due to each activity as detailed in Table 4.12 below.

TABLE 4.12 MANAGEMENT ACTIVITIES, TIER 1 EQUATIONS AND DEFAULT EF FOR DEAD ORGANIC MATTER C POOL CHANGES

Activity	Ecosystem	Tier 1 Equation ¹ and Default EF
----------	-----------	---

Aquaculture ($\Delta C_{\text{DOM-AQ}}$), Salt Production ($\Delta C_{\text{DOM-SP}}$), Extraction ($\Delta C_{\text{DOM-EXT}}$)	Mangrove forest	AQ, SP, EXT-D² : At start of activity, apply Eq. 2.18 or 2.19; $\text{DOM}_{\text{in}} = 0$ and $\text{DOM}_{\text{out}} = \text{DOM}$ oxidized upon disturbance; Table 4.13 AQ, SP : for abandonment phase, apply Eq. 2.18 and 2.20 or 2.19; Table 4.13 AQ : for use phase, $\Delta C = 0$.
	Tidal marsh	$\Delta C = 0$
	Seagrass meadow	
Drainage ($\Delta C_{\text{B-DR}}$)	Mangrove forest	At start of activity, apply Eq. 2.18 and 2.20-2.22 or 2.19; $\text{DOM}_{\text{out}} = \text{DOM}$ oxidized upon disturbance ; Table 4.13
	Tidal marsh	$\Delta C = 0$
	Seagrass meadow	
Harvesting of Aquatic Resources ($\Delta C_{\text{DOM-HARV}}$), Nutrient Enrichment ($\Delta C_{\text{DOM-NUTR}}$), Hydrologic/Sediment Diversion ($\Delta C_{\text{DOM-DIV}}$), Restoration ($\Delta C_{\text{DOM-RES}}$) & Creation ($\Delta C_{\text{DOM-CRE}}$), Other Activities ($\Delta C_{\text{DOM-OTHER}}$)	Mangrove forest	Apply Eq. 2.18 and 2.20 or 2.19; Table 4.13
	Tidal marsh	$\Delta C = 0$
	Seagrass meadow	
Note – ¹ Equations in this section can be found in the 2006 IPCC Guidelines; ² EXT-D = physical removal of DOM pool associated with Extraction activity; EXT-I is not considered as it is only considered to have potentially significant impacts on seagrass meadows		

4.2.2.1 CHOICE OF METHOD

Changes in DOM resulting from management activities that occur in mangrove forests are estimated at the Tier 1 level because they represent potentially large CO₂ emissions to or removals from the atmosphere. Conversions that result in loss of mangrove forest biomass or oxidation of soils can have large implications for CO₂ emissions and removals. For tidal marsh and seagrass meadow, the default assumption is that no changes in C stocks occur at the Tier 1 level of estimation. For significant management activities in mangrove forests and extensive management activities that impact DOM pools in tidal marshes with perennial biomass, Tier 2 and higher estimation methods should be used and these values reported. In these cases, stock changes should be disaggregated relative to climate, vegetation type and salinity where applicable.

TIER 1

Mangrove forest

At Tier 1, for Aquaculture, Salt Production, and Extraction activities, it is assumed that C in DOM stocks is lost as CO₂ emissions in the year of conversion and that no change in DOM C stocks occurs afterward. For these activities, apply either *Gain-Loss* or *Stock-Difference* methods, following Chapter 2, Eq. 2.18 or 2.19 from Volume 4, *2006 IPCC Guidelines*. If applying *Gain-Loss*, apply Eq. 2.18. If applying *Stock-Difference*, apply Eq. 2.19. However, at the start of the activity (construction phase, including extraction), instead of applying Eq. 2.20 to estimate DOM_{in} of Eq. 2.18, $\text{DOM}_{\text{in}} = 0$. For *Stock-Difference*, $\text{DOM}_{\text{t2}} = 0$. In years subsequent to the start of the activity, the Tier 1 assumption is no change in DOM stock.

Tidal marsh and Seagrass meadow

For all management activities, the Tier 1 assumption is no change in DOM stock.

TIERS 2 AND 3

Tier 2 methodologies also employ Equations 2.18 or 2.19. However, country-specific data are encouraged, especially where management activities are expected to have significant impacts on CO₂ emissions and removals. For instance, after a management activity that results in a significant change in aboveground biomass stocks, conversion of aboveground biomass to DOM as litter can be significant. Countries should use in such cases

Second Order Draft

national estimates for litter stocks for mangrove forests and tidal marshes disaggregated by tidal regime, where applicable. Loss estimates of dead wood and litter exports due to tidal advection should also be considered (see Annex 1 regarding C exports). Tier 3 methods may further employ stratification by ecological zone or disturbance regime to reduce uncertainties. Field measurements can further inform and be used to validate model output at Tier 3. For mangroves, Tier 3 methodologies should employ empirical measurements of canopy litter fall and census of downed wood lying on the forest floor. This methodology should involve the establishment of replicate forest plots of at least 5 X 5 m area with monthly census taking of changes in wood and measurement of litter lying within 1 X 1 m traps placed below the canopy but above highest astronomical tide (see Saenger and Snedakar, 1993, for full description of these methods).

4.2.2.2 CHOICE OF EMISSION/REMOVAL FACTORS

TIER 1

Mangrove forest

Countries using a Tier 1 method require data on the default dead wood and litter carbon stocks as defined in Table 4.13. For coastal wetlands that are identified as mangrove forests prior to when the activity took place and activities including aquaculture, salt production, extraction or hydrologic/sediment diversion occur, Tier 1 emission factors for DOM stocks should be used (Table 4.13). The Tier 1 assumption is that carbon stocks in litter and dead wood pools in all non-forested coastal wetlands and in coastal wetlands where activities including nutrient enrichment and harvesting of aquatic resources occur are zero.

Tidal marsh and Seagrass meadow

For all management activities, the Tier 1 emission factor for DOM stock change is zero.

TIER 2

The higher Tier methods described in Chapter 4 for Forest Land and Chapters 5 and 6 on Cropland and Grassland of Volume 4 of the *2006 IPCC Guidelines* will permit better estimates when national data are applied. Additional requirements may arise if the assumption that carbon stocks in dead wood or litter pools of tidal marsh are zero cannot be justified, such as where intensive management activities have occurred.

TIER 3

Tier 3 emission factors include model output and validation and disaggregated data sources.

TABLE 4.13 TIER 1 DEFAULT VALUES FOR LITTER AND DEAD WOOD CARBON STOCKS

Domain	Ecosystem type	Litter carbon stocks of mature forests (tonnes C ha ⁻¹) with 95% CI	Dead wood carbon stocks of mature forests (tonnes C ha ⁻¹) with 95% CI
Tropical/Subtropical	Mangrove forest	0.7 (0-1.3)	10.7 (6.5-14.8)
Source: Litter: Utrera-Lopez and Moreno-Casasola 2008, Liao et al 1990, Chen et al 2008, Richards et al 2011, Ramose-Silva et al 2007, Twilley et al 1986 Dead Wood: Kauffman et al 2011, Donato et al 2012, Allen et al 2000, Steinke et al 1995, Robertson et al 1989, Tam et al 1995, Krauss et al 2005			

4.2.2.3 CHOICE OF ACTIVITY DATA

Choice of activity data follows from guidance provided in the *2006 IPCC Guidelines* in the Forest Land chapter (p. 4.38). Also, refer to Section 4.2.1.3 for guidance relative to specific management activities.

4.2.2.4 CALCULATION STEPS FOR TIER 1

Calculations steps for Tier 1 is presented in the *2006 IPCC Guidelines* and follows from guidance provided in the Forest Land chapter (p. 4.38).

4.2.2.5 UNCERTAINTY ASSESSMENT

Guidance on uncertainty assessment for dead organic matter pools is unchanged from the *2006 IPCC Guidelines* in the Forest Land chapter (p. 4.38). With specific regard to coastal wetlands, uncertainties follow from Tier 1 assumptions about tidal export of C as well as assumptions of only non-woody biomass in tidal marshes.

4.2.3 Soil Carbon

Activities that occur within coastal wetlands can influence organic and mineral stocks of C in soils. The methodological approach follows equation 4.1 which combines the emissions from the activities listed in Table 4.14. Current data on inorganic soil C are not sufficient to provide generic methodologies and so the methodological approach provided here only takes into account the possible emissions and removals associated with changes in organic C of organic and mineral soils.

Management activities that include extraction of saturated soils that are then deposited above the existing water table, or that lead to drainage due to lowering of the water table, result in enhanced soil oxidation. The management activities Aquaculture, Salt Production, Extraction and Drainage result in the most significant effects on the soil carbon pool. These management activities however do not necessarily result in a land-use category change and this consideration needs to be applied when reporting CO₂ emissions. Conversely, restoration or creation of coastal wetlands can result in increased net C accumulation and subsequently net removals of CO₂ from the atmosphere. These activities can also be significant.

Activities associated with either localised or extensive lowering of the water table are often associated with the construction of drainage channels leading to CO₂ fluxes due to oxidation of dissolved organic and particulate organic carbon in the water carried by drainage channels. However, there is currently not enough information to provide emission factors for C exports (see Future Methodological Development).

Second Order Draft

Table 4.14 Management Activities, Tier 1 Approaches and Tier 1 Equations for Soil Carbon Pool Changes		
Activity	Ecosystem	Tier 1 Equation and Default EF
Aquaculture (ΔC_{SO-AQ}), Salt Production (ΔC_{SO-SP})	Mangrove forest	AQ, SP: Apply Eq. 4.2. EF in Tables 4.15-4.18, this section SP: only <i>construction</i> and <i>abandonment</i> phases are considered
	Tidal marsh	
	Seagrass meadow	AQ: Only <i>use</i> phase is considered. Apply Eq. 2.6 ¹ ; Table 4.17, this section,
Extraction (ΔC_{SO-EXT})	Mangrove forest	EXT-D: $\Delta C_{SO-EXT} = A_{c,e} * EF_{CONST,e} * P$ from Eq. 4.2. EF in Tables 4.15 & 4.16, this section
	Tidal marsh	
	Seagrass meadow	If EXT-D , $\Delta C_{SO-EXT} = A_{c,e} * EF_{CONST,e} * P$, Eq. 4.2, Table 4.17, this section; if EXT-I $\Delta C = 0$
Harvesting of Aquatic Resources ($\Delta C_{SO-HARV}$)	Mangrove forest	Apply Eq. 2.6 ¹ ; Table 4.18, this section
	Tidal marsh	$\Delta C = 0$
	Seagrass meadow	
Nutrient Enrichment ($\Delta C_{SO-NUTR}$)	Mangrove forest	NA ²
	Tidal marsh	
	Seagrass meadow	Apply Eq. 2.6 ¹ ; Table 4.19, this section
Hydrologic/Sediment Diversion (ΔC_{SO-DIV})	Mangrove forest	$\Delta C = 0$
	Tidal marsh	
	Seagrass meadow	NA
Drainage (ΔC_{SO-DR})	Mangrove forest	Apply Eq. 4.3; Table 4.20, this section
	Tidal marsh	
	Seagrass meadow	NA
Restoration (ΔC_{SO-RES}) & Creation (ΔC_{SO-CRE})	Mangrove forest	Apply Eq. 4.4; Table 4.21, this section
	Tidal marsh	
	Seagrass meadow	
Other Activities ($\Delta C_{SO-OTHER}$)	Mangrove forest	Apply Eq. 2.6 ¹
	Tidal marsh	
	Seagrass meadow	
Note ¹ Volume 4 of the 2006 IPCC Guidelines		
² no available Tier 1 guidance		

4.2.3.1 AQUACULTURE, SALT PRODUCTION AND EXTRACTION

In aquaculture and salt production, the activity is broken down into three phases — *construction*, *use* and *abandonment* — and emission factors are provided for each of these phases. Extraction effects can take two forms: 1) direct via on-site physical removal of soil (EXT-D) or 2) specifically in seagrass meadows, indirectly via effects of soil deposition and loss of water clarity (EXT-I)

Construction phase and Extraction

In excavation there is only one phase, which is equivalent to the *construction* phase of aquaculture and salt production. During the *construction* phase of shrimp or fish ponds, or for salt production in mangrove forest and tidal marsh ecosystems, soil is excavated to build containing berms within which water is held. Depending on the type of aquaculture (intensive, extensive etc.) and the species stocked in the ponds (shrimp, fish) the soils can be excavated to make ponds of 50 cm to 2.5 m depth (Cruz, 1997; Kungvankij et al. 1986; Wang 1990; Robertson and Phillips 1995). In a similar manner the depth of solar salterns varies between depths of about 0.5 to 2.5m (e.g. Ortiz-Milan 2006, Madkour & Gaballah 2012). Therefore at Tier 1, it is assumed that soil is excavated to 1 m depth and that this is appropriate for aquaculture and salt production in the *construction* phase, as well as extraction. It is assumed that the soil excavated at the year in which these activities start is instantaneously oxidized. Differentiation between organic and mineral soils should also be made when reporting on the *construction* phase of aquaculture and salt production activities and extraction in mangrove forests and tidal marshes. Available data suggest that only direct extraction will impact the soil C pool of seagrass meadows through physical disturbance. Aquaculture in seagrass meadows using fish cages and pens is known to be a significant activity in some coastal areas but requires no soil excavation.

Use phase

In the second or *use* phase of aquaculture and salt production, ponds, pens or cages are stocked or ponds maintained for salt production. In aquaculture, the *use* phase results in CO₂ emissions from nutrient enrichment of the soils due to feeding and growing of stock (Holmer et al. 2003; 2004). For aquaculture, there is currently not enough information to provide emission factors for the *use* phase in tidal marshes and it is assumed that the emission factor is the same as that for, *use*, in mangrove forests. Once aquaculture begins, it is assumed that emissions persist until the ponds (constructed in mangrove forests and tidal marshes) or cages (established in seagrass meadows) are abandoned. The average lifetime of a shrimp pond is variable depending on various factors (management, water quality, and sediment characteristics) and a viability of 7–15 years has been estimated, considering improved management (Flaherty and Karnjakesom 1995, Pa'ez-Osuna 2001). Thus an average lifetime of 10 years is assumed before the activity ends. During the *use* phase, no differentiation is made between organic and mineral soils. For salt production, there is currently not enough information to provide emission factors for the *use* phase.

Abandonment phase

The final phase of aquaculture and salt production is the *abandonment* of the ponds, pens, or cage. In aquaculture the ponds are usually abandoned due to disease or declining water quality (Stevenson et al 1999). During this phase, the area utilized for aquaculture or salt production is assumed to leave the soils in a saturated condition. For abandoned ponds maintained as saturated soils in mangrove forests the emissions are assumed to be the same as for forest clearance until which time vegetation reestablishment begins. On abandonment, the area utilized for aquaculture is generally left unchanged and the ponds can remain unproductive for decades, but sometimes restoration of the local hydrology eventually leads to revegetation. In the *abandonment* phase, solar salterns established in tidal marshes or fish pens and cages constructed in seagrass meadows, there is currently not enough information to provide emission factors for these ecosystem types.

For aquaculture and salt production, the total change in C stocks in organic and mineral soils is estimated using Equation 4.2 below, which combines emissions from soil organic matter decomposition during all phases. Direct extraction of soil (EXT-D) is applied as for the construction term. In seagrass meadows there is the indirect impact of extraction (EXT-I), which affects the soil C pools differently.

EQUATION 4.2

ANNUAL CHANGE IN SOIL C IN COASTAL WETLANDS WITH AQUACULTURE, SALT PRODUCTION ACTIVITIES IN THREE PHASES OF SUB-ACTIVITIES AND EXTRACTION

$$\text{CO}_{2\text{-SOA,e}} = \sum e(A_{\text{CONSTR,e}} * P * \text{EF}_{\text{CONSTR,e}}) + (A_{\text{USE,e}} * \text{EF}_{\text{USE,e}}) + (A_{\text{ABAN,e}} * \text{EF}_{\text{ABAN,e}})]$$

where:

CO_{2SOe} = CO₂ emissions from either organic or mineral soils that are used in Aquaculture or Salt Production occurring in the three phases of *construction*, *use* and *abandonment* or during Extraction and ecosystem type(e) (mangrove forest, tidal marsh and seagrass meadow), unit = tonnes C yr⁻¹

A_e = Total area under aquaculture and salt production activities during the *construction*, *use* and *abandonment* phases (for mangrove forests or tidal marshes), during *use* phase (for seagrass meadows) or

Second Order Draft

during extraction (for mangrove forests, tidal marshes or seagrass meadows) by ecosystem type (e) as specified, unit = ha

EF_{CONSTR_e} = CO₂ emissions from either organic or mineral soils that are excavated for aquaculture or salt production during the *construction* phase (for mangrove forests or tidal marshes) or during extraction (for mangrove forests, tidal marshes or seagrass meadows) by ecosystem type (e) as specified, unit = tonnes C ha⁻¹

P = fraction of area excavated for aquaculture or salt production during the *construction* phase. The same value of P is used during the subsequent phases of *use* and *abandonment*. Tier 1 assumes P=1 for aquaculture or Salt Production, or during *use* phase (for seagrass meadows), dimensionless.

EF_{USE_e} = CO₂ emissions from aquaculture or salt ponds in mangrove forests and tidal marshes or covered by aquaculture cages or pens in seagrass meadows by ecosystem type (e) as specified, unit = tonnes C ha⁻¹ yr⁻¹

EF_{ABAND_e} = CO₂ emissions from saturated soils after abandonment of aquaculture or salt production practices in mangrove forests as specified, unit = tonnes C ha⁻¹ yr⁻¹

a = activity (aquaculture, salt production; extraction method is based on equation provided)

e = ecosystem type (specified ecosystem in which the activity has potentially significant impacts in mangrove forest, tidal marsh, or seagrass meadow)

CHOICE OF METHOD - ORGANIC AND MINERAL SOILS

Tier 1

Mangroves and Tidal Marsh

The method for estimating changes in soil carbon for Tier 1 uses Equation 4.2 for CO₂ emissions from organic and mineral soils. For Tier 1, extraction (EXT-D) is applied in the case of physical removal and CO₂ emissions are reported for either or both ecosystems where this activity occurs. For salt production, emissions are only reported for *construction* (mangrove forest and tidal marsh; Tables 4.15 & 4.16) and *abandonment* (mangrove; Table 4.18). For aquaculture, CO₂ emissions in mangroves and tidal marshes are reported for all three phases, *construction*, *use* and *abandonment* following Equation 4.2. Data pertaining to CO₂ emissions from abandoned aquaculture ponds on tidal marshes were not found and so the Tier 1 CO₂ emissions are assumed to be zero. It is assumed that after 10 years the emissions associated with aquaculture *use* cease as the practise is abandoned.

At Tier 1, the area affected during use and abandonment must remain the same as that during the construction phase and it is assumed that the entire area is affected (P=1).

Seagrass Meadow

For Tier 1, if extraction (direct extraction, EXT-D) is an important management activity, CO₂ emissions are reported. For Tier 1 indirect effects of extraction (EXT-I), the default assumption is that no changes in C stocks occur. For significant aquaculture activities in seagrass meadows, emissions are only reported for the *use* phase. It is assumed that after 10 years the emissions associated with aquaculture *use* cease as the practise is abandoned (Holmer et al., 2003).

Tier 2

During the phases of *construction*, *use*, and *abandonment* of aquaculture and salt ponds or aquaculture cages and pens, the area in which the activities primarily occur may be better approximated as a proportion of the total area identified for Tier 1 estimation. That is, higher Tier methods may define the area of the activity as the area of soil actually excavated to construct the pond or covered by fish pens (i.e. P<1). For significant activities, Tier 2 and higher estimation methods should be used and these values reported. In these cases emissions should be separated relative to climate, ecological zone and salinity where applicable.

At Tier 2 countries should use national data on their particular aquaculture practices to include more specific information on, depth of pond excavated during the construction phase and determine how type of stock and feeding regime during the *use* phase affect CO₂ emissions and removals. Disaggregation of mineral and organic soils during *abandonment*, and consideration of more appropriate differentiation with respect to salinity and climate regime are some higher Tier improvements. For salt production, a Tier 2 level method requires more specific information on depth of pond excavated. For extraction the method countries should use national data to

determine their particular extraction processes and the volume of soil removed as well as taking into account the fate of the excavated soil.

Tier 3

Tier 3 methods can employ models to estimate CO₂ emissions based on the whether the practise involves extensive or intensive farming techniques, the species farmed, stocking density and feeding regime, all of which affect CO₂ emissions during *use* and likely *abandonment* phases. The effect of temperature and salinity on benthic metabolism both seasonally and with climate and ecological zone should also be included. Tier 3 methods using models should be validated with field measurements. In salt production, the CO₂ emissions and removals associated with the use phase would be improved by providing data to quantify these fluxes and how the magnitude of the fluxes vary with the salt pond management practices.

CHOICE OF EMISSION FACTORS - ORGANIC AND MINERAL SOILS

Tier 1

Mangrove forest, Tidal Marsh and Seagrass Meadow

Default Tier 1 emission factors for aquaculture, salt production and extraction in organic and mineral soils are given in Tables 4.54-4.18 and where appropriate, for each phase of the activity, as well as ecosystems affected. These values should be used in conjunction with Equation 4.2 to estimate emissions. Land area must remain the same through all three processes.

TABLE 4.15 EMISSION FACTORS (EF) ASSOCIATED WITH CONSTRUCTION (EF _{CONSTR}) OF AQUACULTURE (AQ), SALT PRODUCTION (SP) AND EXTRACTION (EXT) ON ORGANIC SOILS (TONNES C HA ⁻¹) AT START OF ACTIVITY				
Ecosystem	EF _{CONSTR}	95% CI	range	n
Mangrove forests	471 ¹	454, 490	216 – 727.1	43
Tidal marsh	340 ²	328, 352	221 – 579	35
Seagrass meadow	NA			
¹ Adame et al. (2012), Breithaupt et al. 2012, Chmura et al. 2003, Donato et al. 2011, Kauffinan et al. 2011, Osborne et al. 2011, Vegas-Vilarrúbia et al. 2010 .				
² Anisfeld et al. 1999, Callaway et al. 1996, Callaway et al. 2012, Chmura & Hung 2004, Craft et al. 1988, Craft 2007, Hussein et al. 2004, Kearney & Stevenson 1991, Orson et al. 1998, Markewich et al. 1998, McCaffrey & Thomson 1980.				

TABLE 4.16 EMISSION FACTORS (EF) ASSOCIATED WITH CONSTRUCTION (EF _{CONSTR}) OF AQUACULTURE (AQ), SALT PRODUCTION (SP) AND EXTRACTION (EXT) ON MINERAL SOILS (TONNES C HA ⁻¹) AT START OF ACTIVITY				
Ecosystem	EF _{CONSTR}	95% CI	range	n
Mangrove forests	286 ¹	266, 306	80 - 1376	77
Tidal marshes	226 ²	214, 239	15.6 – 623	82
Seagrass meadow ³	140 ⁴		9.1 – 829	89
¹ Donato et al. 2011, Chmura et al. 2003, Breithaupt et al. 2012, Fujimoto et al. 1999, Adame et al. 2012, Perry & Mendelssohn 2009, Ren et al. 2010, Kauffinan et al. 2011, Ray et al. 2011, Zhang et al. 2012, Khan et al. 2007, Matsui 1998.				
² Cahoon et al. 1996, Callaway et al. 2012, Chmura & Hung 2004, Connor et al. 2001, Craft et al. 1988, Craft 2007, Hatton 1981, Kearney & Stevenson 1991, Livesley & Andrusiak 2012, Loomis & Craft 2010, Morris & Jensen 2003, Oenema & DeLaune 1988, Patrick & DeLaune 1990, Roman et al. 1997, Yu & Chmura 2009.				
³ For EXTRACTION (EXT) only; ⁴ Fourqurean et al 2012				

Second Order Draft

TABLE 4.17 ANNUAL EMISSION FACTORS (EF) ASSOCIATED WITH USE (EF_{USE}) OF AQUACULTURE (AQ) ON ORGANIC AND MINERAL SOILS (TONNES C HA⁻¹ YR⁻¹)

Ecosystem	EF _{USE}	95% CI	range	n
Mangrove forests and Tidal marshes	2.2 ^{1,2}	1.7, 2.7	0.34-5.9	15
Seagrass meadow	9.5 ³	7.8, 11.6	1.4-28.8	22

¹ Alongi 1999b, Alongi et al. 2004, Blackburn et al. 1988, Lovelock et al. 2011;² It is assumed that EF for tidal marsh is the same as that for mangrove forests.³ Holmer et al. 2003, Holmer et al. 2005, Heilskov & Holmer 2001**TABLE 4.18 ANNUAL EMISSION FACTORS (EF) ASSOCIATED WITH ABANDONMENT AFTER AQUACULTURE (AQ) OR SALT PRODUCTION (SP) UNDER SATURATED CONDITIONS (EF_{ABAN-S}) ON ORGANIC AND MINERAL SOILS AND HARVESTING OF AQUATIC RESOURCES (EF_{HARV}) IN MANGROVE FORESTS (TONNES C HA⁻¹ YR⁻¹)**

Ecosystem	EF _{ABAN-S} /EF _{HARV}	95% CI	Range	N
Mangrove forests	1.8 ¹	0.9, 3.5	0.40 – 7.9	4

¹ Alongi et al. (1998), Lovelock et al. (2011), Cahoon et al. (2003). Data from forest clearance assumed to be the same as Aquaculture during the *abandonment* phase.**Tier 2**

Tier 2 could include the use of country specific emission factors for each ecosystem that take into account local climatic factors. For fish and shrimp ponds, the actual area excavated ($P < 1$) and depth to which soil is excavated, should be taken into account as this varies with aquaculture and salt production practices. During the phase, *use*, and the phase, *abandonment* in mangrove forests, the EF could be disaggregated to provide separate values for organic and mineral soils. Furthermore, country-specific data would include emission factors for tidal marshes and seagrass meadows in the *abandonment* phase.

Tier 3

A Tier 3 approach could use models that take into account the time-dependent nature of the CO₂ fluxes over a range of timescales. For example, during the construction phase a pulse of CO₂ efflux from soil directly after mangrove clearing and prior to excavation, followed by a logarithmic decline in CO₂ fluxes over time should be considered.

4.2.3.2 HARVESTING OF AQUATIC RESOURCES

Guidance on annual change in soil carbon stocks on coastal wetlands managed for resource harvest is provided only in the case of deforestation or complete clearing of aboveground biomass in mangrove forests. These emissions result in net CO₂ flux to the water or atmosphere as there are assumed to be no C inputs to soil once aboveground biomass has been removed. It is assumed that mangrove wood harvesting does not affect the height of the water table and that the soils remain saturated.

Selective logging in mangrove forests, harvesting of vegetation in tidal marshes or shellfish in seagrass meadows, are likely to have a low impact on carbon soil stocks and so the Tier 1 C stock change is assumed to be zero. If any of these activities contribute to emissions that represent a key category, then Tier 2 or Tier 3 estimation methodologies are recommended.

CHOICE OF METHOD & EMISSION/REMOVAL FACTORS**Tier 1*****Mangrove forest***

Default Tier 1 emission factor for harvesting of mangrove wood, resulting in forest clearance, is presented in Table 4.17. For harvesting of aquatic resources, the methodology follows Eq.4.2, however only the *abandonment* term is reported.

Tier 2

Tier 2 methods for wood harvesting may take into consideration the species of mangrove being felled and determination of country specific emission factors to disaggregate between organic and mineral soils. It may be possible to disaggregate emission factors for tree clearance and selective logging. At Tier 2, country and ecosystem specific emission factors may be also used to reflect the regional importance of harvesting specific resources.

Tier 3

A Tier 3 approach requires a comprehensive understanding of the processes that lead to CO₂ emissions and removals and the factors that drive these fluxes. The Tier 3 approach will involve country-specific models and/or measurement-based approaches along with disaggregated land-use data to incorporate information on the type of resources being harvested, the nature of the disruption to the soil, tidal regime, and local salinity. Tier 3 modelling would include country-specific emission factors derived or modelled from measurement data encompassing different resource harvesting and the physical methods deployed. The models should capture variation in emission rates driven by soil disruption, seasonal changes in precipitation and temperature and tidal influence.

4.2.3.3 NUTRIENT ENRICHMENT

Guidance on annual change in emissions associated with nutrient enrichment is provided only in the case of mangrove forests and seagrass meadows. Note that data for tidal marshes is insufficient to generate an emission factor for nutrient enrichment and so it is assumed that the emission factor is the same as reported for mangrove forests.

CHOICE OF METHOD AND EMISSION FACTOR**Tier 1**

Mineral and organic soils have been aggregated for default Tier 1 emission factors for Nutrient Enrichment in seagrass meadows and are presented in Table 4.19. The emissions should be quantified by applying Eq. 2.6, Vol. 4 and be used in conjunction with equation 4.2.

TABLE 4.19 ANNUAL EMISSION FACTORS (EF) ASSOCIATED WITH NUTRIENT ENRICHMENT (EF_{NUTR}) ON ORGANIC AND MINERAL SOILS (TONNES C HA⁻¹ YR⁻¹)				
	EF_{NUTR}	95% CI	Range	N
Seagrass Meadow	3.5 ¹	2.7, 4.5	1.9 - 7.2	5
¹ Holmer et al. 2005, Heilskov et al. 2008; No data available for tidal marshes.				

Tier 2 and 3

A Tier 2 approach will involve country-specific emission factors and may take into consideration the source of the nutrient enrichment and their regional importance. Disaggregation of mineral and organic soils should be considered. A Tier 3 approach will use measurement-based approaches along with disaggregated land-use data to incorporate data on variable rates that will better capture variation in emission rates associated nutrient application or discharge rate as well as factors that include consideration of diel fluctuations in temperature, tidal regime, and salinity. If this activity contributes to emissions that contribute to a category that is key, then Tier 2 or Tier 3 estimation methodologies are recommended. If nutrient enrichment is a significant activity constituting managed coastal wetlands in tidal marshes and mangrove forests, Tier 2 or 3 estimation methodologies are also recommended.

4.2.3.4 HYDROLOGIC/SEDIMENT DIVERSION

Annual change in carbon stocks on coastal wetland managed for hydrologic and sediment diversion can occur in mangrove and tidal marsh. The Tier 1 emission factors for soils in all ecosystems as a result of coastal

Second Order Draft

subsidence are also considered to be zero. If this activity contributes to emissions that contribute to a category that is key, then Tier 2 or Tier 3 estimation methodologies are recommended.

4.2.3.5 DRAINAGE

Emissions from drained coastal wetland soils are estimated at Tier 1 for mangrove forests and tidal marshes because they represent potentially large CO₂ emissions to the atmosphere. Annual C losses from both drained mineral and organic soils in mangrove forests and tidal marshes are aggregated at Tier 1 level of estimation using Equation 4.3.

EQUATION 4.3

ANNUAL CHANGE IN CARBON STOCKS IN DRAINED ORGANIC AND MINERAL SOILS

$$CO_{2DR} = \sum_e (A_e \cdot EF_e \cdot F_{PD})$$

where:

CO_{2DR} = CO₂ emissions from organic or mineral soil C through drainage, units = tonnes C yr⁻¹

A = land area under drainage, units = ha

EF_{DR,e} = CO₂ emissions from organic or mineral soil C through drainage ecosystem type (e), units = tonnes C ha⁻¹ yr⁻¹

F_{PD} = fraction of area that is partially drained. Tier 1 assumes F_{PD}=1

TABLE 4.20 ANNUAL EMISSION FACTORS (EF) ASSOCIATED DRAINAGE (EF_{DR}) ON ORGANIC AND MINERAL SOILS (TONNES C HA⁻¹ YR⁻¹)

Ecosystem	EF _{DR}	95% CI	Range	N
Mangrove forests and tidal marshes	7.9 ^{1,2}	6.5, 9.6	1.2 – 43.9	22

¹ Camporese et al. (2008), Deverel & Leighton (2010), Hatala et al. (2012), Howe et al. (2009), Rojstaczer & Deverel (1993).
² Based on aggregated data from tidal marshes and mangrove forests.

CHOICE OF METHOD AND EMISSION/REMOVAL FACTORS

Tier 1 – Organic and Mineral Soils

Mangrove forest and Tidal Marsh

The default Tier 1 assumption for drainage is that organic C is lost linearly over time until the stock is fully oxidised to CO₂, further drainage is initiated, or a new land use has started. Our default assumption is that the water table has been changed to 1 m below the soil surface for organic and mineral soils. If drainage does not result in a land-use change, the inventory compiler is referred to the aggregated emission factor for drained organic and mineral soils in tidal marshes and mangrove forests as reported in Table 4.20.

If drainage results in conversion to a new Land Use category, the inventory compiler is referred to Section 4.3 of this Chapter. It is important under these reporting conditions to retain information about the conversion so that wetlands guidance can be applied if a reversal of drainage conditions occurs.

Tier 2 and Tier 3

The Tier 2 approach should incorporate country specific data on emission for soils disaggregated for mineral and organic soils. Higher Tier methods could take account of differences in the management of the drained coastal wetland, drainage and the fraction of area that is partially drained (F_{PD}) should be determined.

4.2.3.6 RESTORATION AND CREATION

Rewetting saturates the soil, which can lead to colonization by new wetland vegetation, that is, the subsequent natural re-establishment of original vegetation type in mangrove or tidal marsh ecosystems. Alternatively, rewetting of the soil can be followed by active replanting that restores the original vegetation type. In seagrass meadows, soils remain saturated and it is the water quality that must reach a satisfactory level before re-establishment of vegetation or its restoration. The original ecosystem can be either re-established naturally, or by manual replanting. Under these conditions, where vegetation becomes re-established, ecosystem restoration or creation is assumed to occur. Creation occurs where coastal wetlands cannot be verified to have previously occurred but likely occurred given the proximity of the land to the coastal margin. Rehabilitation, a form of creation where the re-establishment of any vegetation type results in soil C sequestration upon maturation (defined as 20 year since establishment).

Guidance for inventories of rewetting and restoration of coastal wetland ecosystems follows several general simplifying assumptions at Tier 1 level of estimation of CO₂ emissions from soil:

1. upon rewetting of previously drained soil, or re-establishment of water quality, creation of biomass by purposeful planting or restoration of biomass is initiated. During this 20 year transient period it is assumed that soil emissions and removals are insignificant, so that rewetting of coastal wetlands results in a Tier 1 EF = 0
2. after the 20 year transient period, restoration results in a change to a coastal wetland ecosystem supporting growth of mangrove, tidal marsh or seagrass biomass, which can then accumulate soil C whereby a Tier 1 soil EF_{RES} is applied. At this time it is assumed that soil C accumulation rates are functionally equivalent to the natural system net soil C accumulation (flux) and the Tier 1 EF_{RES} is applied (Table 4.20) with the same EF applied regardless of soil type (Craft et al 2003, McGlathery et al 2012, Osland et al 2012).

If the re-establishment of functioning vegetation is demonstrated to be unsuccessful during rewetting an, EF=0 is applied for a further 20 year period or until creation or restoration of biomass occurs. For creation it is assumed that once the site has been purposefully planted the EF=0 for soil for a 20 year transient period, by which time functionally equivalent soil C accumulation rates are assumed to occur and EF_{CRE} can be applied. Therefore EF_{RES} is the same as EF_{CRE}

EQUATION 4.4 ANNUAL CHANGE IN SOIL C IN RESTORED AND CREATED COASTAL WETLANDS

$$CO_{2SO-RESe} = (A_{RESe} * EF_{RESe}) \quad CO_{2SO-CREe} = (A_{CREe} * EF_{CREe})$$

where,

CO_{2SO-RESe} = CO₂ emissions associated with rewetted, improved water quality and restored coastal wetlands by ecosystem type (mangrove, tidal marsh and seagrass meadow), units = tonnes C yr⁻¹

CO_{2SO-CREe} = CO₂ emissions associated with rewetted improved water quality and created coastal wetlands by ecosystem type (mangrove, tidal marsh and seagrass meadow), units = tonnes C yr⁻¹

A_{RESe} = Area of soil that has been modified by rewetting, improved water quality and/or restoration, by ecosystem type (e), units = ha

EF_{RESe} = CO₂ emissions from mineral and organic soils that have been modified by restoration, by ecosystem type (e), units = tonnes C ha⁻¹ yr⁻¹

A_{CREe} = Area of soil that has been modified by rewetting and/or creation, by ecosystem type (e), units = ha

EF_{CREe} = CO₂ emissions from mineral and organic soils after creation by ecosystem type (e), units = tonnes C ha⁻¹ yr⁻¹

This method can be applied for lands with either mineral or organic soils. At Tier 1, organic and mineral soils are not differentiated and land area estimates should be based on vegetation classification within the new land category to apply Tier 1 EFs.

CHOICE OF METHOD AND EMISSION/REMOVAL FACTORS

Tier 1

Second Order Draft

Mangrove, Tidal Marsh and Seagrass Meadow

Changes in soil carbon resulting from management activities that occur during rewetting and restoration or creation are estimated at Tier 1 level because they represent potentially large C removals from the atmosphere. During the transitional, rewetting period, the soil EF=0 regardless of the vegetation that is present. An EF=0 is applied until 20 years after vegetation re-establishment. The land can be assumed restored with functioning vegetation unless the definition of a classified vegetation type (as determined by country land classification system) is not met. Once the land enters the new land category, the EF in Table 4.21 can be applied. EF values should be used in conjunction with Equation 4.4 to estimate emissions.

TABLE 4.21 ANNUAL EMISSION FACTORS (EF) ASSOCIATED WITH RESTORATION OR CREATION (EF_{RES/CRE}) ON ORGANIC AND MINERAL SOILS (TONNES C HA⁻¹ YR⁻¹) AFTER 20 YEARS OF VEGETATION REESTABLISHMENT

Ecosystem	EF _{RES/CRE}	95% CI		range	n
Mangrove	1.62 ¹	0.73	3.63	0.10 – 10.2	69
Tidal marsh	0.91 ²	0.82	1.02	0.05 – 4.65	66
Seagrass meadow	0.43 ³	0.28	0.67	0.09 – 1.12	6

¹ Breithaupt et al. 2012, Chmura et al. 2003, Fujimoto et al. 1999, Ren et al. 2010.

² Anisfeld et al 1999, Cahoon et al. 1996, Callaway et al 1996, Callaway et al 1997, Callaway et al 1998, Callaway et al 1999, Callaway et al. 2012, Chmura & Hung 2003, Hatton 1981, Craft 2007, Kearney & Stevenson 1991, Markewich et al. 1998, Oenema & DeLaune 1988, Orson et al 1998, Patrick & DeLaune 1990, Roman et al 1997.

³ Mateo & Romero 1997, Serrano et al. 2012.

Tier 2 and 3

For key categories, Tier 2 and higher estimation methods should be used and these values reported. Soil types, organic and mineral, should be disaggregated in order to apply suitable country-specific factors. Country-specific emission factors applied during the rewetting of the lands and the transition period should be used under Tier 2 methods, and the inventory compilers are encouraged to estimate the transient time from rewetted to restored wetlands. Under the tier 3 method, the land use prior to rewetting, climate and vegetation type should be taken into account. A comprehensive understanding and representation of the dynamics CO₂ gas emission factors, based on field measurement is involved in Tier 3. A Tier 3 approach would also use models that take into account the time-dependent nature of the CO₂ fluxes over a range of timescales and the effects of sea-level rise on soil C sequestration rates (Morris et al., 2012) or other dynamics (Craft 2001).

4.2.3.7 OTHER ACTIVITIES

This category is retained to draw attention to the fact that all significant activities may not be covered in this guidance. It is good practice to consider other activities which may, based on national circumstances, have significant C emissions or removals associated with that management activity. A generic method is provided in Table 4.13.

4.2.3.8 CHOICE OF ACTIVITY DATA

See guidance provided in Section 4.2.1.3.

4.2.3.9 CALCULATION STEPS FOR TIER 1

The following summarizes steps for estimating change in carbon stocks in soils for Land Remaining in a Land-use Category where managed coastal wetlands occur.

Step 1: Organize data into inventory time periods based on the years in which activity data were collected (e.g., 1990 and 1995, 1995 and 2000, etc.).

Step 2: Determine the area of coastal wetland in the last year of the inventory time period.

Step 3: Determine the activities to be included in the assessment and their representative areas. Area data should be obtained using the methods described in Chapter 3 of the *2006 IPCC Guidelines*.

Step 4: Determine the activity categories by stratifying representative areas into soil type and previous coastal wetland vegetation type.

Step 5: For each activity category, assign the appropriate emission factor (EF) for annual losses of CO₂.

Step 6: Estimate total emissions by summing the product area (A) multiplied by the emission factor (EF) for all activity categories.

Step 7: Repeat for additional inventory periods.

Example: The following example shows the calculations for aggregate areas of soil carbon change in coastal wetlands. There is a 1Mha area of coastal wetlands. At the beginning of the inventory time period (1990 in this example) the total area of coastal wetlands was 500,000 ha. From this total 200,000 ha was identified as unmanaged and 300,000 ha was actively managed. The individual areas relevant to activities included in the inventory are; 50,000 ha of area under aquaculture with organic soils in previous mangrove forest (with 50% under *construction* and 50% under *use*); 10,000 ha of area under aquaculture with mineral soils in previous tidal marsh (with 90% under *use* and 10% *abandoned*); 50,000 ha of area under fish cages in seagrass meadows (100% under *use*); 100,000 ha of area under salt production with mineral soils in previous tidal marsh (with 100% under *construction*); and 90,000 ha of area under hydrologic diversion. The CO₂ emissions for the inventory time period are calculated as (50,000 ha • 0.5 • 471 tonnes C ha⁻¹) + (50,000 ha • 0.5 • 2.2 tonnes C ha⁻¹) + (10,000 ha • 0.9 • 2.2 tonnes C ha⁻¹) + (10,000 ha • 0.1 • 7.9 tonnes C ha⁻¹) + (50,000 ha • 9.5 tonnes C ha⁻¹) + (100,000 ha • 340 tonnes C ha⁻¹) + (90,000 ha • 0 tonnes C ha⁻¹) = 46,332,700 tonnes C for 1990. Repeat for additional inventory periods.

4.2.3.10 UNCERTAINTY ASSESSMENT

Few studies have addressed the questions of emission factors specifically and few reports are available to give specifics of activities which we expect to vary geographically. Thus, derivation of emission factors was based upon expert knowledge and application of data from reports of biomass and production of vegetation, the amount of carbon in living biomass and the amount of carbon held in soils, and how easily it is decomposed. There is uncertainty in the time, depth of soil affected, the rate of CO₂ loss from dead vegetation and soils and the C gained during wetland creation or restoration.

4.2.4 Non-CO₂ emissions

This section provides methods for estimating the emissions of N₂O and CH₄ from coastal wetlands under different management activities.

N₂O is an intermediate product of both nitrification (oxidation of ammonium by nitrifying microbes under oxic or aerobic conditions) and denitrification (reduction of nitrite and nitrate by denitrifying microbes under anoxic conditions). The flux of N₂O thus is controlled by oxygen availability and tidal influence, but also by the availability of N substrates (Purvaja and Ramesh, 2001; Kreuzwieser et al., 2003; van den Heuvel et al., 2009; Moseman-Valtierra et al. 2011). Therefore, management activities that change hydrology or N availability in sediment, e.g. N management or rewetting of wetlands, could lead to changes in N₂O emissions from coastal wetlands.

In reduced and anoxic environments microbial decomposition of the organic matter may produce CH₄. However, flooding seawater contains sulfate and microbial reduction of sulfate to sulfide will generally occur before methanogens produce CH₄. A strong inverse relationship between CH₄ emission and salinity of mangrove soils was reported by Purvaja and Ramesh (2001) and later work by Poffenbarger et al. (2011) showed that polyhaline tidal marshes (salinity >18) had significantly lower CH₄ emissions than oligohaline (salinity 0.5-5) and mesohaline (salinity 5-18) marshes. The depth distribution of methanogenesis in tidal marsh soils is closely related to the methanogenic substrate utilization (Parkes et al. 2012).

Second Order Draft

Aquaculture has been shown to lead to changes in N₂O emissions from coastal wetlands. Shrimp and fish cultivation increases nutrients loads in culture ponds. The body of research is growing, and sufficient for developing emission factors for non-CO₂ emissions from this activity but is relatively limited. Coastal wetlands subject to intensive carbon and nutrient loading may also be sources of CH₄ emissions, however, only when salinity <18 are present. In aquaculture, anaerobic conditions are not appropriate for maximized fish production and are therefore assumed negligible, however, they may increase over time.

If soil conditions prior to rewetting were aerobic, then it is likely that CH₄ emissions will increase where tidal water salinities are <18 ppt. If tidal waters rewetting coastal wetlands are carbon- or nutrient-enriched, then wetlands with salinities >18 ppt also are likely to be sources of CH₄ emissions. One of the main controlling factors in N₂O production is the availability of inorganic N in the soil. If, prior to rewetting, the area of coastal wetland was subject to increased N supply (e.g. in fertilized land) it may be a source of N₂O emissions during the rewetting. However, experimental studies have shown that the substrate N is rapidly depleted (Moseman-Valtierra et al. 2011), thus N₂O is likely to be negligible after the N is depleted and no continuous N is supplied.

N₂O emissions from managed terrestrial soils are covered in Chapter 11, Volume 4 of the 2006 IPCC Guidelines, which estimates N₂O emissions due to N additions to soils (e.g. synthetic or organic fertilizers, deposited manure, crop residues, sewage sludge), or of mineralization of N in soil organic matter following management of organic soils or cultivation on mineral soils (e.g., Forest Land/Grassland/Settlements converted to Cropland). For organic soils, these emission factors have been updated for these specific land uses and can be found in Chapter 2 of this Supplement Table 4.21 presents the equations and emission factors for managed coastal wetland. The most significant activities contributing to non-CO₂ emissions from managed coastal wetlands are Aquaculture and specific cases of nutrient enrichment, such as aquaculture effluent (indirect N₂O emissions from aquaculture use) for which there are available data Non-CO₂ emissions are assumed equal to zero except in the cases of Aquaculture effects on N₂O emissions and nutrient enrichment effects on CH₄ emissions caused by aquaculture and agricultural run-off in coastal wetlands.

TABLE 4.22 ACTIVITIES AND TIER 1 ASSUMPTIONS FOR NON-CO₂ EMISSIONS FOR THE ECOSYSTEMS AFFECTED

Activity	Ecosystem	Non-CO ₂ gas/sub-activity	Tier 1 Equation
Aquaculture (EF _{AQ}) ¹	Mangrove	CH ₄ : EF=0	NA
		N ₂ O: EF=based on fish production or N fed or nutrient-enriched EF	Apply Eq. 4.5; Table 4.23
	Tidal Marsh	CH ₄ : EF=0	NA
		N ₂ O: EF=based on fish production or N fed	Apply Eq. 4.5; Table 4.23
	Seagrass Meadow	CH ₄ : EF=0 N ₂ O: EF=0	NA
Salt Production (EF _{SP}), Extraction (EF _{EXT}), Hydrologic/Sediment Diversion (EF _{DIV}), Harvesting of Aquatic Resources (EF _{HARV}), Drainage (EF _{DR})	Mangrove	CH ₄ : EF=0 N ₂ O: EF=0	NA
	Tidal Marsh	CH ₄ : EF=0 N ₂ O: EF=0	
	Seagrass Meadow	NA	
Nutrient Enrichment (EF _{NUTR})	Mangrove	CH ₄ : EF=nutrient-enriched wetland (agricultural run-off including aquaculture effluent)	Apply Eq. 2.6 ² ; table 4.24, this section
		N ₂ O: Run-off from terrestrial soil N application	Chapter 11 ²
		N ₂ O: Enrichment from aquaculture effluent	Apply Eq. 2.6 ² ; table 4.25, this section

	Tidal Marsh	CH ₄ : EF=nutrient-enriched wetland (agricultural run-off including aquaculture effluent)	Apply Eq. 2.6 ² ; table 4.24, this section
		N ₂ O: Run-off from terrestrial soil N application	Chapter 11 ²
		N ₂ O: Enrichment from aquaculture effluent	Apply Eq. 2.6 ² ; table 4.25, this section
	Seagrass Meadow	CH ₄ : EF=0	NA
		N ₂ O: Run-off from terrestrial soil N application	Chapter 11 ²
		N ₂ O: Enrichment from aquaculture effluent	NA
Restoration (EF _{RES}) and Creation (EF _{CRE})	Mangrove	CH ₄ : EF=unmanaged wetland N ₂ O: EF=0	Apply Eq. 2.6 ² ; table 4.26, this section
	Tidal Marsh	CH ₄ : EF=unmanaged wetland N ₂ O: EF=0	
	Seagrass Meadow	NA	NA
Other Activities (EF _{OTHER})	Mangrove	NA	NA
	Tidal Marsh		
	Seagrass Meadow		

Note

¹for aquaculture, only the *use* phase is considered; EF for phases of *construction* and *abandonment* are assumed negligible

²Volume 4 of the 2006 IPCC Guidelines

4.2.4.1 AQUACULTURE

CHOICE OF METHOD AND EMISSION FACTORS

Seitzinger et al. (2000) estimated that one-third of global anthropogenic N₂O emissions are from aquatic ecosystems, but there has been limited research aimed at quantifying the contribution of aquaculture to the global budget. N₂O is emitted from aquaculture systems as a by-product of the conversion of ammonia (contained in fish urea) to nitrate through nitrification and nitrate to N₂ gas through denitrification (Hu et al. 2012). Hu et al. (2012) estimates that 1.8% of the nitrogen fed to aquaculture systems is emitted as N₂O, and that 1.7 g N₂O-N is emitted per kg fish produced in an aquaculture system. These are applied only during the *use* phase of aquaculture. In the *construction* and *abandonment* phases, non-CO₂ emissions are assumed negligible and EF=0.

Tier 1

Methane

The Tier 1 method estimates the CH₄ emissions from aquaculture ponds are assumed negligible and reported as 0.

Nitrous Oxide

N₂O emissions from aquaculture ponds can be estimated based on fish production or the amount of N feed in the aquaculture activity. The default Emission Factors for these two methods are provided in Table 4.23. The emission estimation follows a modified form of Eq. 11.1 from Chapter 11, Volume 4 of the 2006 IPCC Guidelines and is presented here in (Eq. 4.5).

Second Order Draft

EQUATION 4.5
DIRECT N₂O EMISSIONS FROM AQUACULTURE USE

$$N_2O-N_{AQ} = F_F * EF_F \text{ OR } F_I * EF_I$$

Where:

N_2O-N_{AQ} = annual direct N₂O-N emissions from Aquaculture use, kg N₂O-N yr⁻¹

F_F = annual fish production, kg fish yr⁻¹

EF_F = emissions factor for N₂O emissions from fish produced, kg N₂O-N (kg fish produced)⁻¹

F_I = annual N amount of fish feed applied

EF_I = emission factor for N₂O emissions from N of fish feed applied, kg N₂O-N (kg N fed)⁻¹

Table 4.23 Emission factors for N₂O emission from aquaculture in coastal wetlands

Wetland	Default EF	Uncertainty Range	Reference
Aquaculture	0.003 kg N ₂ O per kg fish produced per yr		Hu et al. 2012
	0.028 kg N ₂ O per kg N fed per yr		

Tiers 2 and 3

Under Tier 2 method, country specific emission factors for these two gases need to be applied. N₂O emissions from aquaculture ponds could also be estimated using the country specific emission rate from the water if such data is available. In terms of CH₄, Tier 2 or 3 methods could include change in CH₄ emissions with extended use, based on age of the pond and intensity of use. A comprehensive understanding and representation of the dynamics based on direct field measurements or models is involved under Tier 3 method, which estimates emission factors considering the category of aquaculture (fish species or feed stuff), aquaculture use intensity, and impact of environmental factors e.g. climate zone, season, and salinity.

4.2.4.2 SALT PRODUCTION, EXTRACTION, HYDROLOGIC/SEDIMENT DIVERSION, HARVESTING OF AQUATIC RESOURCES, DRAINAGE

As was noted in Table 4.22, CH₄ and N₂O emissions from the management activities of salt production, extraction, hydrologic/sediment diversion and harvesting of aquatic resources are assumed to be zero. However, under higher Tiers, emissions associated with these activities should be reported using country-specific data.

As was noted in Table 4.22, CH₄ and N₂O emissions from the management activities of drainage are assumed to be zero. However, if drainage is a significant activity within the Land Use category under which it is reported, it is good practice to apply Tier 2 and 3 estimation levels following generic methodologies (Eq. 2.6, 2006 IPCC Guidelines) and using country-specific data.

4.2.4.3 NUTRIENT ENRICHMENT

CHOICE OF METHOD AND EMISSION/REMOVAL FACTORS

Methane and Nitrous Oxide - Mangrove and Tidal Marshes

Tier 1

For the Tier 1 method, the default emission factors are given in Tables 4.24 and 4.25. The default method to estimate non-CO₂ gas emissions from coastal wetlands is to multiply wetland area by the gas emission rates. For nutrient enrichment from agricultural run-off in the coastal zone, only CH₄ emissions are reported. N₂O emissions for this activity are covered in Chapter 11 of the *2006 IPCC Guidelines*. For nutrient enrichment from aquaculture effluent coming from aquaculture production both CH₄ and N₂O emissions are reported as there is currently no guidance in *2006 IPCC Guidelines* on non-CO₂ emissions from this impact of aquaculture activities. Data were not available on impact of aquaculture effluent on seagrass meadows.

Calculation of CH₄ emissions follows the same method applied in the case of agricultural run-off to the coastal zone and aquaculture effluent within the coastal zone (Table 4.24). For calculation of N₂O emissions associated with agricultural run-off from terrestrial sources that comes from either direct or indirect N application to terrestrial soils the inventory compiler is referred to Chapter 11 of the *2006 IPCC Guidelines*. The following assumption is applied under Tier 1 estimation: vegetation of coastal wetlands is not a source of non-CO₂ gases and the emission factor represents only fluxes directly from the soil to atmosphere.

A second method for estimating indirect N₂O emissions from nutrient enrichment resulting from aquaculture effluent uses a modified form of Eq. 11.9 and 11.10 for effluent effects from aquaculture. The leaching factor can be applied and the EF from estuaries following this method. This method is recommended when absence of data on land area impacted by aquaculture effluent prevents use of that approach (see Chapter 11, Volume 4 of the *2006 IPCC Guidelines*).

TABLE 4.24 EMISSION FACTORS FOR CH₄ EMISSIONS FROM NUTRIENT ENRICHMENT (AGRICULTURAL RUN-OFF AND AQUACULTURE EFFLUENT) (TONNES CH₄ HA⁻¹ YR⁻¹)

Ecosystem	EF	95% CI	range	n
Mangrove forest and Tidal marsh (oligohaline - polyhaline)	1.13 ¹	0.68, 1.18	0.015 – 27.83	18

¹ Mukhopadhyay et al., 2002, Kreuzwieser et al., 2003, Alongi et al., 2005, Biswas et al., 2007, Chauhan et al., 2008, Krithika et al., 2008, Liikanen et al., 2009, Allen et al., 2010, Chen et al., 2010, Tong et al., 2010, Adams et al., 2011.

TABLE 4.25 EMISSIONS FACTORS FOR N₂O EMISSIONS FROM NUTRIENT ENRICHMENT (AQUACULTURE EFFLUENT) (KG N₂O HA⁻¹ YR⁻¹)

Ecosystem	EF	95% CI	range	n
Mangrove forest and Tidal marsh (oligohaline - polyhaline)	3.7 ¹	0.19, 5.4	-4.6 – 66.3	17

¹ Corredor et al., 1999, Bauza et al., 2002, Kreuzwieser et al., 2003, Alongi et al., 2005, Krithika et al., 2008, Liikanen et al., 2009, Chen et al., 2010, Adams et al., 2011, Allen et al., 2011, Chen et al., 2012.

Tier 2 and 3

Tier 2 methods are used where countries have their country-specific emission factors and substantial national data. The Tier 2 method allows the inventory compiler to adjust the assumptions applied to the Tier 1 method. It is a good practice to use the representative data of climate zones and vegetation type from published literature, e.g. papers, reports and books, and estimate the non-CO₂ gases emissions from wetland plants, where applicable.

Tier 3 methods involve a comprehensive understanding and representation of the dynamics of non-CO₂ gas emission factors, taking account parameters such as salinity, season, N load, tidal elevation and tidal cycle (high tide vs. low tide). Field research should be carefully carried out at representative sites for empirical gas fluxes. Countries are encouraged to setup their own methodology or modelling exercises based on the N input rates for estimation of emission factors.

4.2.4.4 RESTORATION AND CREATION

The degree of water saturation is a key factor controlling CH₄ and N₂O emissions. Rewetting of coastal wetland soils shifts microbial decomposition from aerobic to anaerobic conditions, increasing the potential for CH₄

Second Order Draft

emissions. If water used to rewet coastal wetlands increases N supplies through ground water or tidal water, a new source of N₂O emissions is possible.

Guidance for inventories of rewetting and restoration or creation of coastal wetlands follows several general simplifying assumptions at Tier 1 level of estimation.

- (i) The default assumption is that water introduced for rewetting, by whatever means, is ambient water free of excess nutrients.
- (ii) Once rewetting occurs, the EF equals 0 for N₂O during a 20 year transition period is the default transition period unless the re-establishment of functioning vegetation is demonstrated to be unsuccessful.
- (iii) Once rewetting occurs, the EF for CH₄ in Table 4.27 should be applied during a 20 year transition period.
- (iv) After the 20 year transition period which begins with the reestablishment of vegetation, the EF remains the same for both N₂O and CH₄ while the land continues to be subject to the management activity of restored or created, regardless of the land-use category that is applied.

In the case of rewetting of lands that had been previously been in agricultural (or any other drained) land-use or category, for example, the prior land use is not considered at Tier 1 as any increase in emissions from rewetting has not only been shown to be a transient effect and data are insufficient to assign an EF to capture this effect. The water quality should be reevaluated to determine whether the EF for nutrient-enriched wetlands should continue to be applied. If water applied in the rewetting activity is enriched in nutrients and the restoration activity contributes to a land-use category or pool that is key, Tier 2 and 3 methods should be applied.

CHOICE OF METHOD AND EMISSION FACTORS

Tier 1

Mangrove and Tidal Marsh

The Tier 1 method in this section estimates the non-CO₂ emissions without considering the land use prior to rewetting. The Tier 1 default assumption is that the lands are rewetted with uncontaminated water. Thus, the emission factor for N₂O = 0 and for CH₄, emission factors are found in Table 4.26. For mangrove forests, the EF for CH₄ is zero. Default Tier 1 emission factors for rewetting and restoration or creation of coastal wetlands with organic and mineral soils are given in Table 4.21. These values should be used in conjunction with equation 4.2 to estimate emissions and take into account the ambient soil salinity and vegetation type. For tidal marshes, the CH₄ EF is based on salinity (which is controlled by factors such as proximity to ocean water, the supply of freshwater through precipitation and fluvial inputs, and evapotranspiration).

TABLE 4.26 EMISSION FACTORS FOR CH₄ FROM UNMANAGED COASTAL WETLANDS FOR TIER 1 ESTIMATION OF REWETTING AND RESTORATION

Wetland Type	Salinity type	Salinity (ppt)	Default EF (tonnes CH ₄ -C ha ⁻¹ y ⁻¹)	EF Range (tonnes CH ₄ -C ha ⁻¹ y ⁻¹)	Error (95%CI)	Reference
Tidal Marsh	Tidal fresh	0.5-5	1.12	0.03-4.04	±80%	Poffenbarger et al. 2011 ¹
	Oligohaline/Mesohaline	5-18	0.28	0.02-0.24		Poffenbarger et al. 2011 ¹
	Polyhaline (>18ppt)	>18	0	0-0.04	±90%	Poffenbarger et al 2011 ¹
Mangrove	Saline	>18	0		±90%	Page and Dal 2010

Note ¹Review paper

Tier 2 and Tier 3

Country-specific emission factors applied during the rewetting of the lands and the transition period should be used under Tier 2 methods, and the inventory compilers are encouraged to estimate the EF for the land during the transition to the new land category for the rewetted lands. Under the tier 3 method, the land use prior to rewetting, climate and vegetation type should be taken into account. A comprehensive understanding and representation of the dynamics of non-CO₂ gas emission factors, based on field measurement, is involved in Tier 3. Tier 2 could include the use of country specific emission factors for each ecosystem that take into account local climatic factors, salinity, and species composition of the vegetation. The quality of the water should also be considered in the case of seagrass restoration.

In the case of rewetting of lands that had been previously been in agricultural (or any other drained) land-use or category, for example, the prior land use is not considered at Tier 1 because data are insufficient to assign an EF to capture this transient effect. The emissions of N₂O and CH₄ are considered to also depend on the nutrient/organic matter available in the tidal water. The water quality should be reevaluated to determine whether the EF for nutrient-enriched wetlands should continue to be applied. If water applied in the rewetting activity is enriched in nutrients and the restoration activity contributes to a land-use category or pool that is key, Tier 2 and 3 methods should be applied.

Tier 2 and 3 should also consider the effect of vegetation as plants can act as conduit for gas exchange between the soil and atmosphere.

4.2.4.5 CHOICE OF ACTIVITY DATA

Aquaculture ponds

For aquaculture, this is the fish production, the amount of feed, and amount of N in feed. These data can be obtained from FAO (<http://www.fao.org/fishery/statistics/global-aquaculture-production/en>). To estimate non-CO₂ gas emissions from aquaculture activities, the area under cultivation must be determined either by on-site measurement or from aerial photographs. Also required is the rate of nitrogen input as feed, which can usually be obtained from the pond operator, and the sediment type beneath the cages or at the bottom of the pond. For additional guidance, see Section 4.2.1.3

Nutrient enrichment

To estimate CH₄ emissions, the area receiving agricultural runoff must be determined by visual inspection of the maximum ingress of tidal waters within the wetland. For Tier 2 and 3 methods, the rate of N input should be determined although this may be very difficult and may involve knowledge of the tidal prism and N concentrations in the waters in question. For nutrient enrichment, this is the area of nutrient-affected land. We can expect to find nutrient-affected systems downstream of major urban centers, municipal sewage treatment facilities, and watersheds where a large portion of the land is under intensive agricultural use. For additional guidance, see Section 4.2.1.3.

Restoration/Creation

To estimate non-CO₂ gas emissions, the area restored, created and/or rewetted must be determined by visual inspection of pre- and post-impact stages. The type of wetland and the salinity regime must be measured. For additional guidance, see Section 4.2.1.3.

4.2.4.6 UNCERTAINTY ASSESSMENT

There have been few empirical measurements upon which to base emission factors and the factors identified for incorporation in Tier 2 and 3 will reflect variability. Uncertainties in non-CO₂ emissions associated with aquaculture can be reduced greatly by better estimation of fish production, quality of N in feed, area of ponds relative to total area under aquaculture use. For additional guidance, see Section 4.2.1.3.

4.3 CONVERSION FROM A LAND USE CATEGORY THAT INCLUDES COASTAL WETLANDS

This section references methodology provided in Section 4.2 to estimate impacts of soil drainage, extraction or other activity when lands are classified under a land-use category other than Wetland. Examples of when classification of land into a different land-use category may occur are given in Box 4.1. In these cases, the

inventory compiler is directed to the guidance in those chapters of Volume 4 of the *2006 IPCC Guidelines* (Chapter 5-Croplands, Chapter 8-Settlements) but applying the assumptions and default EFs provided here.

4.3.1 CO₂ emissions and removals

4.3.1.1 BIOMASS, DOM AND SOIL CARBON

Permanent drainage under agriculture use or conversion to Settlements are cases in which a land-use conversion may be considered to have occurred. For biomass and DOM, see sections 4.2.1 and 4.2.2 unless land is converted to lands with dry soils in which case follow guidance in *2006 IPCC Guidelines* (following flow chart provided in Chapter 1). For soil carbon, refer to Chapter 2, this supplement for drained organic soils and the respective Chapters in the *2006 IPCC Guidelines* for land use conversion to lands with mineral soil. Regardless of how the land is classified, supplementary guidance provided in this Chapter should be followed where management activities where coastal wetlands are involved.

4.3.2 Non-CO₂ emissions

This section provides guidelines for estimating non-CO₂ gas emissions from coastal wetlands, especially mangroves and tidal marshes being drained for conversion to other dry land uses. The land use following coastal wetland drainage (e.g., agriculture, settlement, forestry), will have a major impact on emissions of non-CO₂ greenhouse gases from soils. Refer to the respective chapter depending on end Land Use in the *2006 IPCC Guidelines*. For guidance on drainage of organic soils and conversion to another IPCC Land-use category, refer to the guidance provided in Chapter 2, this supplement.

4.4 CONVERSION TO A LAND USE CATEGORY THAT INCLUDES COASTAL WETLANDS

4.4.1 CO₂ emissions and removals

This section includes management activities that result in rewetting and restoration or creation of coastal wetland ecosystems and subsequent classification as Wetland. Land that is exposed to rewetting can previously have sustained mangroves or tidal marshes, but the previous land-use is characterised by (A) unsaturated soils due to sediment infill to a height above the water table, (B) drainage of the original ecosystem, (C) construction of dikes or levees that prevent tidal flooding, or (D) emplacement of culverts that restrict flow of tidal waters.

Methods for biomass and DOM pools follow guidance provided in Sections 4.2.1 and 4.2.2. Non-CO₂ emissions are covered in section 4.4.4 and follow a similar approach with EFs for rewetting applied during the transitional land-use and EFs for restoration or creation applied when functioning vegetation has become re-established. For example, an aquaculture facility that is classified as Settlement or Cropland and restored to Wetlands would involve a change in IPCC land use category.

Guidance for inventories of rewetting and restoration of coastal wetland ecosystems follows similar general simplifying assumptions at Tier 1 level of estimation of C emissions from soil (as presented in Section 4.2.3): 1) upon rewetting, EF=0 for a 20 year *transition* period (as opposed to *transient* period in the case of Land Remaining in a Land-use Category). The default transition period is 20 years unless the re-establishment of functioning vegetation is demonstrated to be unsuccessful, 2) after conversion to the new (restored or created coastal) land use category, a new EF is applied. The new EF is equivalent to the natural system net soil C accumulation (flux). Regardless of the land category, reestablishment of functioning vegetation is assumed to occur after a 20 year transition period (in the case of a land-use change to Wetlands) and a new EF is applied for soils (Table 4.21) following general guidance in Section 4.2.3.

4.4.2 Non-CO₂ emissions

This section provides guidelines for estimating non-CO₂ gas emissions from coastal wetlands, especially mangroves and tidal marshes, undergoing management activities that result in rewetting and restoration or creation of coastal wetland ecosystems. These activities will have a major impact on emissions of non-CO₂ greenhouse gases from soils. Apply methods and default EF provided in section 4.2.4.5.

4.5 COMPLETENESS, TIMES SERIES CONSISTENCY, QUALITY ASSURANCE AND QUALITY CONTROL

4.5.1 Completeness

Completeness is a requirement for greenhouse gas inventories, and it is *good practice* to address all wetland carbon gain and losses. For completeness, it is *good practice* to include all carbon pools and non-CO₂ gases. The wetland area used for calculation for different carbon pools must be the same, and emissions from organic and mineral soils on mineral soils should be estimated. Country-specific information should be incorporated into higher tier methodologies. A complete accounting of emissions and removals of CO₂ associated with for Land Remaining in a Land-use category, Land Converted to another Land-use category in which the land that is converted is a coastal wetland, and Conversion to a Land-Use category that includes coastal wetlands regardless of how it is classified. It is *good practice* that all losses from biomass carbon pools that result in transfers to dead organic matter pools are first accounted for as changes to biomass carbon stocks.

4.5.2 Developing a consistent time series

It is good practice to develop a consistent time series of inventories of anthropogenic emissions and removals of greenhouse gases using the guidance in Chapter 7 in this volume. Achieving time series data may require extrapolation or interpolation from longer time series data or from long term trends, as few long-term data are available for most coastal wetlands.

Consistent accounting over time of wetland areas included in biomass and soil C emissions and removals inventory requires that activity data be stratified by the common definitions of wetland type/ soil type. Wetlands subject to land-use change must not be lost or double-counted due to accounting errors resulting from inconsistent stratification of wetland types/ soil types. Ideally, the same protocol should be applied consistently every year in the time series, at the same level of disaggregation and where country-specific data are used, it is *good practice* to use the same values and methods for equivalent calculations throughout the time series.

New values should be included if the inventory capacity and information and data sources improve over time. It is *good practice* in these circumstances to consistently recalculate the earlier emissions and removals. Other changes during the time series need to be consistent to take account of new data or methods and their consistency with the earlier data. It is *good practice* to recalculate the entire time series of data if the default values are changed; changes in wetland types need to be tracked for long periods of time.

It is *good practice* to use the same model parameter values for the entire time series and to recalculate the entire dataset if one of more of the parameters has changed. Failure to do so may result in either under-or over-estimates of the true changes in carbon and non-CO₂ gas emissions or removals.

4.5.3 Quality assurance and quality control

Different levels of precision and accuracy, and as a result, bias will invariably apply to a number of the values used to assess greenhouse gas inventories. Estimates are influenced by the quality and consistency of data and information available as well as knowledge gaps, all of which will vary among countries. Depending on the tier level used, estimates will be affected by different sources and degrees of error, such as sampling error.

It is *good practice* to execute quality control checks through Quality Assurance (QA) and Quality Control (QC) procedures as detailed in Chapter 7, and review the emission estimation procedures by experts. Additional quality control checks as outlined in Chapter 7 and quality assurance procedures may also be applicable. This is especially so if higher tier methods are used. It is *good practice* to supplement the general QA/QC related to data

Second Order Draft

processing, handling, and reporting and documenting, with source-specific procedures. QA/QC procedures should be documented separately for Land Remaining in a Land-use category, Land Converted to another Land-use category in which the land that is converted is a coastal wetland, and Conversion to a Land-Use category that includes coastal wetlands regardless of how it is classified.

Organizations and institutions which collect the data are responsible for reviewing data collection methods and all aspects of the data handling and analysis procedures, and ensure that they are done correctly, and are complete and consistent over time. It is important to document all procedures and processes as it enables reviewers to identify inaccuracy, gaps and to suggest improvements. Transparency is most important in order to ensure consistency and clarity of the processes and procedures over time.

All data should be checked against other reliable sources of information that are independent. Any differences or discrepancies must be documented, and consistency must be applied to total areas involved in the inventory to ensure that wetland area are neither 'created' or 'lost' overtime. When using country-specific data, the inventory compiler should compare these data to the IPCC default values or the Emissions Factor Database (EFDB) and detail any differences. These country-specific data must be of high quality, adequately described, and documented.

If factors are based on direct measurements (i.e., soil C content) the inventory agency should review the measurements to ensure that they are representative of the actual range of environmental conditions. It is *good practice* to review and, if necessary, revise the default assumptions and to compare model estimates with field measurements and other data sources.

4.5.4 Reporting and documentation

General requirements for reporting and documentation are set out in Chapter 7. It is *good practice* to archive and document all data and information applied to produce the national emissions/removals inventory. Definitions of all carbon pools should be included in the inventory, including evidence that these definitions have been applied consistently over time.

Documentation is necessary for demonstrating transparency, completeness, consistency of all data and methods for interpolating between samples, methods and years, and for recalculating and avoidance of possible double accounting or 'loss' of C inventory. Regardless of Tier methodology used, explanations are required for decisions regarding choice of methodology, approaches and use of default or other data. This is necessary to facilitate examination by independent third parties; inventories should include summaries of approaches and methods used and references to data sources so that the reported emissions estimates are transparent and can be retraced or recalculated.

All data sources, including default values, must be quoted. The scientific basis for any country-specific data and methods must be completely described and justified, as well as describing sources and magnitudes of uncertainty. This is especially so for any large-scale estimates as in these cases the statistical procedures should be described and well as the level of uncertainty.

Differences between years in emissions should be explained and the possible reasons for these differences documented as much as possible.

4.6 FUTURE METHODOLOGICAL DEVELOPMENT

4.6.1 C export

The amount of dissolved and particulate carbon potentially available for export is highly variable among coastal wetlands, depending on a large number of factors such as: net primary productivity, tidal range, the ratio of wetland to watershed area, lateral trapping of tidal water, the presence of high salinity plugs in the tropical dry season, total wetland area, frequency of storms, amount of precipitation, and volume of water exchange. Each ecosystem is unique; some wetlands export DOC but import POC, others import DOC and POC but export DIC, while other systems import or export all forms of dissolved and particulate carbon. The direction of net exchange also usually varies within the same estuary with change in season.

Accurate estimation of tidal exchange in a particular wetland is not a straightforward process. Many workers have provided rough estimates by multiplying carbon concentrations suspended in wetland creeks and waterways by the tidal range multiplied by the creek/waterway cross-sectional area. Estimates derived from such simple calculations are invalid and misleading for a number of reasons, including the inherent assumption that there are differences in carbon concentrations between ebb and flood tide stages and that the tidal prism is symmetrical. In fact, carbon concentrations in many wetland waters do not show significant differences between tides. Further, tides in most wetlands are asymmetrical, characterized by a pronounced asymmetry between ebb and flood tides with the ebb most often being of shorter duration but with stronger current velocity than the flood tide. Also, tidal velocities vary across a waterway with faster surface current velocities mid-stream than those just above the creekbed or proximal to the wetland.

For these reasons, it is not possible to make simple generalizations regarding total carbon export from mangroves, seagrasses or tidal marshes and, in fact, comparatively few such measurements have been made properly. The correct method would be to measure water volume and velocity over entire tidal cycles over several seasons in relation to position in the water-column to derive an overall annual estimate of average water flow by volume. This involves fairly complex instrument measurements and sophisticated mathematical modelling as well as extensive and expensive repetitive measurements of dissolved and particulate carbon concentrations. For mangroves, net exchange of carbon has been properly measured in only twelve systems, with no clear exchange patterns among locations, although it does appear that most mangroves export POC as litter but with rates ranging widely from 0.1-27.7 mol C m⁻²yr⁻¹ (Alongi, 2009). This export equates globally to only about 10% of total carbon fixed by trees; respiration to the atmosphere is by far the largest loss of C to the atmosphere. Such appears to be the case for tidal marshes (Chmura et al., 1993) and subtidal seagrass beds (Fourqurean et al., 2012)

References

- Aber JS, F Pavri, S. Aber. 2012 Wetland Environments: A Global Perspective. Wiley-Blackwell.
- Abohassan, R.A.A., C.A. Okia, J.G. Agea, J.M. Kimondo and M.M. McDonald, 2012. Perennial Biomass Production in Arid Mangrove Systems on the Red Sea Coast of Saudi Arabia, Environmental Research Journal 6(1), pp. 22-31.
- Adame, M. F., Reef, R., Herrera-Silveira, J. A., and Lovelock, C. E. (2012). Sensitivity of dissolved organic carbon exchange and sediment bacteria to water quality in mangrove forests. *Hydrobiologia*, **691**:239–253.
- Adams, C.A., Andrews, J.E., Jickells, T. (2011). Nitrous oxide and methane fluxes vs. carbon, nitrogen and phosphorous burial in new intertidal and saltmarsh sediments. *Science of the Total Environment* (in press)
- Agostini, S., Pergent, G., & Marchand, B. (2003). Growth and primary production of *Cymodocea nodosa* in a coastal lagoon. *Aquatic botany*, 76(3), 185-193.
- Aioi, K. (1980). Seasonal change in the standing crop of eelgrass (*Zostera marina* L.) in Odawa Bay, central Japan. *Aquat. Bot.* 8: 343-354.
- Aioi, K., & Pollard, P. C. (1993). Biomass, leaf growth and loss rate of the seagrass *Syringodium isoetifolium* on Dravuni Island, Fiji. *Aquatic botany*, 46(3), 283-292.
- Aioi, K., Mukai, H., Koike, I., Ohtsu, M., & Hattori, A. (1981). Growth and organic production of eelgrass (*Zostera marina* L.) in temperate waters of the Pacific coast of Japan II. Growth analysis in winter. *Aquatic Botany*, 10, 175-182.
- Ajonina, G. N. 2008. Inventory and Modelling Mangrove Forest Stand Dynamics Following Different Levels of Wood Exploitation Pressures in the Douala-Edea Atlantic Coast of Cameroon, Central Africa (Doctoral dissertation, Faculty of Forest and Environmental Sciences, Albert-Ludwigs-Universität).
- Alberts, J. J., Filip, Z., Price, M. T., Williams, D. J., & Williams, M. C. 1988. Elemental composition, stable carbon isotope ratios and spectrophotometric properties of humic substances occurring in a salt marsh estuary. *Organic geochemistry*, 12(5), 455-467.
- Allen, J. A., Ewel, K. C., Keeland, B. D., Tara, T., & Smith, T. J. 2000. Downed wood in Micronesian mangrove forests. *Wetlands*, 20(1), 169-176.

Second Order Draft

- 1573 Allen D.E., Dalal R.C., Rennenberg H., Meyer R.L., Reeves S. and Schmidt S. 2007. Spatial and temporal
1574 variation of nitrous oxide and methane flux between subtropical mangrove soils and the atmosphere. *Soil*
1575 *Biology and Biochemistry* 39: 622–631.
- 1576 Allen, D., Dalal, RC, Rennenberg, H, Schmidt, S. 2011. Seasonal variation in nitrous oxide and methane
1577 emissions from subtropical estuary and coastal mangrove sediments, Australia. *Plant Biology* 13: 126-133.
- 1578 Alongi, DM. 2010. *The Energetics of Mangrove Forests*. Springer Science.
- 1579 Alongi, D.M. 2011. Early growth responses of mangroves to different levels of nitrogen and phosphorus supply.
1580 *Journal of Experimental Marine Biology & Ecology* 397: 85-93.
- 1581 Alongi, DM, BF Clough, P. Dixon, F Tirendi. 2003. Nutrient partitioning and storage in arid-zone forests of the
1582 mangroves *Rhizophora stylosa* and *Avicennia marina*. *Trees* 17: 51-60.
- 1583 Alongi, D. M., Sasekumar A, Chong CV, Pfizner J, Trott LA, Tirendi F, Dixon P, Brunskill GJ (2004). Sediment
1584 accumulation and organic material flux in a managed mangrove ecosystem: estimates of land-ocean-
1585 atmosphere exchange in peninsular Malaysia. *Mar Geol*, **208**: 383-402.
- 1586 Alongi, D. M., Tirendi, F., & Trott, L. A. (1999b). Rates and pathways of benthic mineralization in extensive
1587 shrimp ponds of the Mekong delta, Vietnam. *Aquaculture*, 175(3), 269-292.
- 1588 Alongi, D.M., McKinnon, A.D., Rachmansyah, Brinkman, R., Trott, L.A., Muana, Undu, M.C., 2009. The fate
1589 of organic matter derived from small-scale fish cage aquaculture in coastal waters of Sulawesi and Sumatra,
1590 Indonesia. *Aquaculture* 295: 60-75.
- 1591 Alongi, D.M. and 7 others. 2001. Organic carbon accumulation and metabolic pathways in sediments of
1592 mangrove forests in southern Thailand. *Mar. Geol.* 179: 85-103
- 1593 Alongi, DM, Pfizner, J, Trott, LA, Tirendi, F, Dixon, P, Klumpp, DW. 2005. Rapid sediment accumulation and
1594 microbial mineralization in forests of the mangrove *Kandelia candel* in the Jiulongjiang Estuary, China.
1595 *Estuarine Coastal and Shelf Science* 63: 605-618.
- 1596 Alongi, D.M., Wattayakorn, G., Tirendi, F. and Dixon, P. 2004. Nutrient capital in different aged forests of the
1597 mangrove *Rhizophora apiculata*. *Botanica Marina* 47:116-124.
- 1598 Amarasinghe, M. D., Balasubramaniam, S. (1992). Net primary productivity of two mangrove forest stands on
1599 the northwestern coast of Sri Lanka. *Hydrobiologia* (247:), 37-47.
- 1600 Anisfeld, S. C., Tobin, M. J., & Benoit, G. (1999). Sedimentation rates in flow-restricted and restored salt
1601 marshes in Long Island Sound. *Estuaries and Coasts*, 22(2), 231-244.,
- 1602 Apostolaki ET, Holmer M, Marbà N, Karakassis I. 2010. Metabolic imbalance in coastal vegetated (*Posidonia*
1603 *oceanica*) and unvegetated benthic ecosystems. *Ecosystems* 13:459–471
- 1604 Apostolaki, ET Holmer, M Marba, N (Karakassis, I 2011 Reduced carbon sequestration in a Mediterranean
1605 seagrass (*Posidonia oceanica*) ecosystem impacted by fish farming AQUACULTURE ENVIRONMENT
1606 INTERACTIONS Volume: 2 Issue: 1 Pages: 49-59Armentano, T. V. and Menges, E. S. 1986. Patterns of
1607 change in the carbon balance of organic soil wetlands of the temperate zone. , *Journal of Ecology* 74:755–
1608 774.
- 1609 Asmus, R. M., Sprung, M., & Asmus, H. (2000). Nutrient fluxes in intertidal communities of a South European
1610 lagoon (Ria Formosa)–similarities and differences with a northern Wadden Sea bay (Sylt-Rømø Bay).
1611 *Hydrobiologia*, 436(1), 217-235.
- 1612 Auby, I., & Labourg, P. J. (1996). Seasonal dynamics of *Zostera noltii* hornem. In the bay of arcachon (France).
1613 *Journal of Sea Research*, 35(4), 269-277.
- 1614 Augier, H., Calvert, H., Wollaston, E., & Santimone, M. (1982). A comparison of the C, H, N, protein and amino
1615 acid composition of *Posidonia australis* Hook. f. with that of *Posidonia oceanica* (L.) delile and several other
1616 marine phanerogams. *Aquatic Botany*, 12, 69-80.
- 1617 Bandeira, S. O. (1997). Dynamics, biomass and total rhizome length of the seagrass *Thalassodendron ciliatum* at
1618 Inhaca Island, Mozambique. *Plant Ecology*, 130(2), 133-141.

- 1619 Bandeira, S. O. (2002). Leaf production rates of *Thalassodendron ciliatum* from rocky and sandy habitats.
1620 Aquatic botany, 72(1), 13-24.
- 1621 Bay, D. (1984). A field study of the growth dynamics and productivity of *Posidonia oceanica* (L.) delile in Calvi
1622 Bay, Corsica. Aquatic Botany, 20(1), 43-64.
- 1623 Bauza, J., Morell, J.M, Corredor, J.E. (2002). Biogeochemistry of nitrous oxide production in the red mangrove
1624 (*Rhizophora mangle*) forest soils. Estuarine, Coastal and Shelf Science 55: 697-704.
- 1625 Benito, I., Onaindia, M. (1991) Biomass and aboveground production of 4 angiosperms in Cantabrian (N. Spain)
1626 salt marshes. Vegetatio 96, 165-175.
- 1627 Birch, J. B., and J. L. Cooley. 1982. Production and standing crop patterns of giant cutgrass [*Zizaniopsis*
1628 *miliacea*] in a freshwater tidal marsh. *Oecologia* 52: 230-235.
- 1629 Blackburn, T. H., Lund, B. A., & Krom, M. D. (1988). C- and N-mineralization in the sediments of
1630 earthen marine fishponds. Mar. Ecol. Prog. Ser., 44: 221-227.
- 1631 Boon, P. I. (1986). Nitrogen pools in seagrass beds of *Cymodocea serrulata* and *Zostera capricorni* of Moreton
1632 Bay, Australia. Aquatic botany, 25, 1-19.
- 1633 Borum, J., Pedersen, O., Greve, T. M., Francovich, T. A., Zieman, J. C., Fourqurean, J. W., and Madden, C. J.
1634 (2005). The potential role of plant oxygen and sulfide dynamics in die-off events of the tropical seagrass,
1635 *Thalassia testudinum*. *Journal of Ecology* **93**, 148-158. doi:10.1111/ J.1365-2745.2004.00943.
- 1636 Bose AN, Ghosh SN, Yang CT, and Mitra A. 1991. Coastal Aquaculture Engineering. Routledge, Chapman and
1637 Hall, Inc. NY, NY, 371 pp.
- 1638 Bouchard, V., Lefeuvre, J.C. (2000) Primary production and macro-detritus dynamics in a European salt marsh:
1639 carbon and nitrogen budgets. Aquatic Botany 67, 23-42.
- 1640 Bouillon S, AV Borges, E Castaneda-Moya, K Diele, T. Dittmar, NC Duke, E Kristensen, SY Lee, C Marchand,
1641 JJ Middelburg, VH Rivera-Monroy, TJ Smith III, RR Twilley. 2007. Mangrove production and carbon sinks:
1642 A revision of global budget estimates. Global Biogeochem Cycles 22: GB2013, doi:
1643 10.1029/2007GB0030S2
- 1644 Boyer, K.E., Callaway, J.C., Zedler, J.B. (2000) Evaluating the progress of restored cordgrass (*Spartina foliosa*)
1645 marshes: Belowground biomass and tissue nitrogen. Estuaries 23, 711-721.
- 1646 Breithaupt, J. L., Smoak, J. M., Smith III, T. J., Sanders, C. J., & Hoare, A. 2012. Organic carbon burial rates in
1647 mangrove sediments: Strengthening the global budget. Global Biogeochemical Cycles, 26(3), GB3011.
- 1648 Briggs, S. V. (1977). Estimates of biomass in a temperate mangrove community. Australian Journal of
1649 Ecology(2), 369-373.
- 1650 Brouns, J. J. (1985). A comparison of the annual production and biomass in three monospecific stands of the
1651 seagrass *Thalassia hemprichii* (Ehrenb.) aschers. Aquatic botany, 23(2), 149-175.
- 1652 Brouns, J. J. (1987b). Aspects of production and biomass of four seagrass species (Cymodoceaceae) from Papua
1653 New Guinea. Aquatic botany, 27(4), 333-362.
- 1654 Brouns, J. J., & Heijs, F. M. (1986). Production and biomass of the seagrass *Enhalus acoroides* (Lf) Royle and its
1655 epiphytes. Aquatic botany, 25, 21-45.
- 1656 Brun, F. G., van Zetten, E., Cacabelos, E., & Bouma, T. J. (2009). Role of two contrasting ecosystem engineers
1657 (*Zostera noltii* and *Cymodocea nodosa*) on the food intake rate of *Cerastoderma edule*. Helgoland Marine
1658 Research, 63(1), 19-25.
- 1659 Bulthuis, D. A., & Woelkerling, W. J. (1983). Biomass accumulation and shading effects of epiphytes on leaves
1660 of the seagrass, *Heterozostera tasmanica*, in Victoria, Australia. Aquatic Botany, 16(2), 137-148.
- 1661 Burd, A. B., & Dunton, K. H. (2001). Field verification of a light-driven model of biomass changes in the
1662 seagrass *Halodule wrightii*. Marine Ecology Progress Series, 209, 85-98.
- 1663 Cabaço, S., Santos, R., & Duarte, C. M. (2008). The impact of sediment burial and erosion on seagrasses: A
1664 review. *Estuarine, Coastal and Shelf Science*, 79(3), 354-366.

Second Order Draft

- 1665 Cabaco, S., Santos, R. & Sprung, M. 2012. Population dynamics and production of the seagrass *Zostera noltii* in
1666 colonizing versus established meadows. *Marine Ecology*, 33, 280–289.
- 1667 Cahoon, D. R., Hensel, P., Rybczyk, J., McKee, K. L., Proffitt, C. E. & Perez, B. C. (2003) Mass tree mortality
1668 leads to mangrove peat collapse at Bay Islands, Honduras after Hurricane Mitch. *Journal of Ecology*, **91**:
1669 1093-1105.
- 1670 Cahoon, D. R., Lynch, J. C., & Knaus, R. M. (1996). Improved Cryogenic Coring Device for Sampling Wetland
1671 Soils: RESEARCH METHOD PAPER. *Journal of Sedimentary Research*, 66(5).
- 1672 Callaway, J. C., DeLaune, R. D., & Patrick Jr, W. H. (1996). Chernobyl 137Cs used to determine sediment
1673 accretion rates at selected northern European coastal wetlands. *Limnology and Oceanography*, 444-450.
- 1674 Callaway, J. C., Borgnis, E. L., Turner, R. E., & Milan, C. S. (2012). Carbon Sequestration and Sediment
1675 Accretion in San Francisco Bay Tidal Wetlands. *Estuaries and Coasts*, 1-19.
- 1676 Calleja, M. L., Barrón, C., Hale, J. A., Frazer, T. K., & Duarte, C. M. (2006). Light regulation of benthic sulfate
1677 reduction rates mediated by seagrass (*Thalassia testudinum*) metabolism. *Estuaries and coasts*, 29(6), 1255-
1678 1264.
- 1679 Camacho, L. D., Gevaña, D. T., Carandang, A. P., Camacho, S. C., Combalicer, E. A., Rebugio, L. L., & Youn,
1680 Y. C. 2011. Tree biomass and carbon stock of a community-managed mangrove forest in Bohol, Philippines.
1681 *Forest Science and Technology*, 7(4), 161-167.
- 1682 Cambridge, M. L., & Hocking, P. J. (1997). Annual primary production and nutrient dynamics of the seagrasses
1683 *Posidonia sinuosa* and *Posidonia australis* in south-western Australia. *Aquatic Botany*, 59(3), 277-295.
- 1684 Carruthers, T. J. (1994). Leaf production, canopy structure and light climate of a density manipulated
1685 *Amphibolis griffithii* meadow. University of Western Australia.
- 1686 Cartaxana, P., Catarino, F. (1997) Allocation of nitrogen and carbon in an estuarine salt marsh in Portugal.
1687 *Journal of Coastal Conservation* 3, 27-34.
- 1688 Cebrián, J., Duarte, C. M., Marbà, N., & Enriquez, S. (1997). Magnitude and fate of the production of four
1689 cooccurring western Mediterranean seagrass species. *Marine Ecology Progress Series*, 155, 29-44.
- 1690 Cebrián, J., Pedersen, M. F., Kroeger, K. D., & Valiela, I. (2000). Fate of production of the seagrass *Cymodocea*
1691 *nodosa* in different stages of meadow formation. *Marine Ecology Progress Series*, 204, 119-130.
- 1692 Chauhan, R., Ramanathan, A.L., Adhya, TK. 2008. Assessment of methane and nitrous oxide flux from
1693 mangroves along the Eastern coast of India. *Geofluids* 8: 321-332.
- 1694 Chave, J., C. Andalo, S. Brown, M.A. Cairns, J.Q. Chambers, D. Eamus, H. Folster, F. Fromard, N. Higuchi, T.
1695 Kira, J.P. Lescure, B.W. Nelson, H. Ogawa, H. Puig, B. Riera, T. Yamakura. 2005. Tree allometry and
1696 improved estimation of carbon stocks and balance in tropical forests. *Oecologia* 145: 87-99.
- 1697 Chen G.C., Tam N.F.Y. and Ye Y. 2010. Summer fluxes of atmospheric greenhouse gases N₂O, CH₄ and CO₂
1698 from mangrove soil in South China. *Science of the Total Environment* 408: 2761–2767.
- 1699 Chen G.C., Ye Y., Lu C.Y. 2008. Seasonal variability of leaf litter removal by crabs in a *Kandelia candel*
1700 mangrove forest in Jiulongjiang Estuary, China. *Estuarine, Coastal and Shelf Science* 79: 701-706.
- 1701 Cheng X., Peng R. Chen J., Luo Y., Zhang Q., An S., Chen J., and Li B. 2007. CH₄ and N₂O emissions from
1702 *Spartina alterniflora* and *Phragmites australis* in experimental mesocosms, *Chemosphere* 68:420-427.
- 1703 Chmura, G.L. and Hung, G. 2004. Controls on salt marsh accretion: a test in salt marshes of Eastern Canadian.
1704 *Estuaries* 27:70-81.
- 1705 Chmura G.L., Anisfeld S., Cahoon D. and Lynch J. 2003. Global carbon sequestration in tidal, saline wetland
1706 soils. *Global Biogeochemical Cycles* 17: 1-12.
- 1707 Christensen, B. 1978. Biomass and primary production of *Rhizophora apiculata* B1 in a mangrove in southern
1708 Thailand. *Aquatic Botany* 4: 43-52.
- 1709 Coles, R. G., Long, W. L., Watson, R. A., & Derbyshire, K. J. (1993). Distribution of seagrasses, and their fish
1710 and penaeid prawn communities, in Cairns Harbour, a tropical estuary, Northern Queensland, Australia.
1711 *Marine and Freshwater Research*, 44(1), 193-210.

- Collier, C. J., Lavery, P. S., Ralph, P. J., & Masini, R. J. (2009). Shade-induced response and recovery of the seagrass *Posidonia sinuosa*. *Journal of Experimental Marine Biology and Ecology*, 370(1), 89-103.
- Camporese, M., Putti, M., Salandin, P., Teatini, P. 2008. Spatial variability of CO₂ efflux in a drained cropped peatland south of Venice, Italy. *Journal of Geophysical Research* 113, doi:10.1029/2008JG000786.
- Connor, R., Chmura, G.L. (2000) Dynamics of above- and belowground organic matter in a high latitude macrotidal saltmarsh. *Marine Ecology-Progress Series* 204, 101-110.
- Connor, R. F., Chmura, G. L., & Beecher, C. B. (2001). Carbon accumulation in Bay of Fundy salt marshes' Implications for. *Global Biogeochemical Cycles*, 15(4), 943-954.
- Coronado-Molina, C., Day, J., Reyes, E., Perez, C. (2004). Standing crop and aboveground biomass partitioning of dwarf mangrove forest in Taylor river Slough, Florida. *Wetlands Ecology and Management* (12:), 157-164.
- Corredor, JE, Morell, JM, Bauza, J. 1999. Atmospheric nitrous oxide fluxes from mangrove sediments. *Marine Pollution Bulletin* 38: 473-478.
- Craft, C. (2007). Freshwater input structures soil properties, vertical accretion, and nutrient accumulation of Georgia and US tidal marshes. *Limnology and oceanography*, 1220-1230.
- Craft, C. B., S. W. Broome, et al. (1988). "Nitrogen, phosphorus, and organic carbon pools in natural and transplanted marsh soils." *Estuaries* 11(4): 272-280.
- Craft, C., Megonigal, J. Patrick, Broome, S., Cornell, J., Freese, R., Stevenson, J., Zheng, L. and Sacco, J. 2003. Pace of ecosystem development of constructed *Spartina alterniflora* in marshes. *Ecological Applications*, 13: 1414-1432
- Craig, N.J., Turner, R.E. and Day Jr., J.W. 1(1979). Land loss in coastal Louisiana (U.S.A.). *Environmental Management* 3: 133-144.
- Cranford, P.J., Gordon, D.C., Jarvis, C.M. (1989) Measurement of cordgrass, *Spartina alterniflora*, production in a macrotidal estuary, Bay of Fundy. *Estuaries* 12, 27-34.
- Cruz, P. S. 1997. Aquaculture feed and fertilizer resource atlas of the Philippines. FAO.
- Curco, A., Ibanez, C., Day, J.W., Prat, N. (2002) Net primary production and decomposition of salt marshes of the Ebre delta (Catalonia, Spain). *Estuaries* 25, 309-324.
- Cyrus, D. P., MacKay, C. F., & Weerts, S. P. 2008. Intrusion of beach-disposed dredger spoil into the Mhlathuze Estuary, South Africa, and its impact on *Zostera capensis*. *African Journal of Aquatic Science*, 33(3), 223-231.
- da Cunha Lana, P., Guiss, C., Disaro, S.T. (1991) Seasonal variation of biomass and production dynamics for above- and belowground components of a *Spartina alterniflora* marsh in the euhaline sector of Paranagua Bay (SE Brazil). *Estuarine Coastal and Shelf Science* 32, 231-241.
- Da Silva J.F., Duck R.W., Catarino J.B. 2004. Seagrasses and sediment response to changing physical forcing in a coastal lagoon. *Hydrology and Earth System Sciences* 8: 151-159.
- Daby, D. (2003). Some quantitative aspects of seagrass ecology in a coastal lagoon of Mauritius. *Marine Biology*, 142(1), 193-203.
- Dahl, T.E. 2011. Status and trends of wetlands in the conterminous United States 2004 to 2009. U.S. Department of the Interior; Fish and Wildlife Service, Washington, D.C. 108 pp.
- Dai M. and others. 2008. Nitrification and inorganic nitrogen distribution in a large perturbed river/estuarine system: the Pearl River Estuary, China. *Biogeosciences* 5: 1545-1585.
- Dame, R.F., Kenny, P.D. (1986) Variability of *Spartina alterniflora* primary production in the euhaline North Inlet estuary. *Marine Ecology-Progress Series* 32, 71-80.
- Darby, FQ and Turner, RE. 2008. Below- and aboveground *Spartina alterniflora* production in a Louisiana salt marsh. *Estuaries and Coasts* 31: 223-231.
- de Boer, W. F. (2000). Biomass dynamics of seagrasses and the role of mangrove and seagrass vegetation as different nutrient sources for an intertidal ecosystem. *Aquatic Botany*, 66(3), 225-239.

Second Order Draft

- 1758 De Iongh, H. H., Wenno, B. J., & Meelis, E. (1995). Seagrass distribution and seasonal biomass changes in
1759 relation to dugong grazing in the Moluccas, East Indonesia. *Aquatic Botany*, 50(1), 1-19.
- 1760 Deegan LA, Johnson DS, Warren RS, Peterson BJ, Fleeger JW, Fagherazzi S, and Wollheim WM 2012 Coastal
1761 Eutrophication as a Driver of Salt Marsh Loss. *Nature* 490: 388-392.
- 1762 De la Cruz, A. A., & Banaag, J. F. 1967. The ecology of a small mangrove patch in Matabungkay Beach,
1763 Batangas Province. *Natural and Applied Science Bulletin*, 20(4), 486-494.
- 1764 Dennison, W. C., & Alberte, R. S. (1985). Role of daily light period in the depth distribution of *Zostera*
1765 *marina*(eelgrass). *Marine ecology progress series*. Oldendorf, 25(1), 51-61.
- 1766 Deverel, S. J. and Leighton, D. A. 2010. Historic, recent and future subsidence, Sacramento-San Joaquin Delta,
1767 California, USA. *San Francisco Estuary and Watershed Science* 8: 1-23.
- 1768 Devereux, R., Yates, D. F., Aukamp, J., Quarles, R. L., Jordan, S. J., Stanley, R. S., & Eldridge, P. M. (2011).
1769 Interactions of *Thalassia testudinum* and sediment biogeochemistry in Santa Rosa Sound, NW Florida.
1770 *Marine Biology Research*, 7(4), 317-331.
- 1771 Dillon, R.C.A. 1971. A comparative study of the primary productivity of estuarine phytoplankton and
1772 macrobenthic plants. Ph. D. thesis, Univ. North Carolina, Chapel Hill, NC.
- 1773 Dittmar, T., R. J. Lara. 2001. Molecular evidence for lignin degradation in sulphate-reducing mangrove
1774 sediments. *Geochim. Cosmochim Acta* 65: 1417-1428
- 1775 Dixon, L. K., & Leverone, J. R. (1995). Light requirements of *Thalassia testudinum* in Tampa Bay, Florida: final
1776 report.
- 1777 Donato D, JB Kauffman, D Murdiyarso, S Kumianto, M Stidham, M Kanninen. 2011. Mangroves among the
1778 most carbon-rich forests in the tropics. *Nature Geoscience* 4: 293-297.
- 1779 Donato D., Kauffman, J. B., MacKenzie, R. A., Ainsworth, A., Pfleeger, A. Z. 2012. Whole-island carbon stocks
1780 in the tropical Pacific: Implications for mangrove conservation and upland restoration. *Journal of*
1781 *Environmental Management* 97:89-96.
- 1782 Dos Santos, V. M., Mathesona, F. E., Pilditch, C. A. & Elger, A. 2012. Is black swan grazing a threat to seagrass?
1783 Indications from an observational study in New Zealand. *Aquatic Botany* 100:41– 50.
- 1784 Doumlele, D.G. (1981) Primary production and seasonal aspects of emergent plants in a tidal freshwater marsh.
1785 *Estuaries* 4, 139-142.
- 1786 Duarte C.M. 1991 Seagrass depth limits. *Aquatic Botany*, 40, 363-377
- 1787 Duarte, C. M., Martínez, R., & Barrón, C. (2002). Biomass, production and rhizome growth near the northern
1788 limit of seagrass (*Zostera marina*) distribution. *Aquatic botany*, 72(2), 183-189.
- 1789 Dumbauld B.R., Ruesink J.L., Rumrill S.S. 2009 The ecological role of bivalve shellfish aquaculture in the
1790 estuarine environment: A review with application to oyster and clam culture in West Coast (USA) estuaries.
1791 *Aquaculture* 290: 196–223.
- 1792 Dunton, K. H. (1990). Production ecology of *Ruppia maritima* L. sl and *Halodule wrightii* Aschers, in two
1793 subtropical estuaries. *Journal of Experimental Marine Biology and Ecology*, 143(3), 147-164.
- 1794 Dunton, K. H. (1994). Seasonal growth and biomass of the subtropical seagrass *Halodule wrightii* in relation to
1795 continuous measurements of underwater irradiance. *Marine Biology*, 120(3), 479-489.
- 1796 Dunton, K. H. (1996). Photosynthetic production and biomass of the subtropical seagrass *Halodule wrightii*
1797 along an estuarine gradient. *Estuaries and Coasts*, 19(2), 436-447.
- 1798 Ellison, A.M., Bertness, M.D., Miller, T. (1986) Seasonal patterns in the belowground biomass of *Spartina*
1799 *alterniflora* (Gramineae) across a tidal gradient. *American Journal of Botany* 73, 1548-1554.
- 1800 Ellison, AM and Farnsworth, EJ. 1996. Anthropogenic disturbance of Caribbean mangrove ecosystems: past
1801 impacts, present trends, and future predictions. *Biotropica* 28: 549-565.
- 1802 Elsey-Quirk, T., Seliskar, D.M., Sommerfield, C.K., Gallagher, J.L. (2011) Salt Marsh Carbon Pool Distribution
1803 in a Mid-Atlantic Lagoon, USA: Sea Level Rise Implications. *Wetlands* 31, 87-99.

- 1804 Erftemeijer, P. L. (1994). Differences in nutrient concentrations and resources between seagrass communities on
1805 carbonate and terrigenous sediments in South Sulawesi, Indonesia. *Bulletin of Marine Science*, 54(2), 403-
1806 419.
- 1807 Erftemeijer P.L.A. and Lewis III, R.R. 2006 Environmental impacts of dredging on seagrasses: A review.
1808 *Marine Pollution Bulletin* 52:1553–1572.
- 1809 Erftemeijer PLA Shuail DA 2012 Seagrass habitats in the Arabian Gulf: distribution, tolerance thresholds and
1810 threats
1811 AQUATIC ECOSYSTEM HEALTH & MANAGEMENT Volume: 15 Special Issue: SI Supplement:
1812 1 Pages: 73-83
- 1813 Fatoyinbo, T. E., Simard, M., Washington-Allen, R. A., & Shugart, H. H. 2008. Landscape-scale extent, height,
1814 biomass, and carbon estimation of Mozambique's mangrove forests with Landsat ETM+ and Shuttle Radar
1815 Topography Mission elevation data. *J. Geophys. Res.*, 113, G02S06.
- 1816 Fatoyinbo, T. E., & Simard, M. 2013. Height and biomass of mangroves in Africa from ICESat/GLAS and
1817 SRTM. *International Journal of Remote Sensing*, 34(2), 668-681.
- 1818 Filip Z, Alberts JJ, Chesire MV, Goodman BA and Bacon JR. 1988. Comparison of salt marsh humic acid with
1819 humic-like substances from the indigenous plant species *Spartina alterniflora*. *Sci. Tot. Environ.* 71: 157-172.
- 1820 Flaherty, M. and Karnjanakesorn, C. 1995. Marine shrimp aquaculture and natural-resource degradation in
1821 Thailand. *Environmental Management* 19: 27-37.
- 1822 Food and Agriculture Organisation of the United Nations (FAO). 2007. The State of the World Fisheries and
1823 Aquaculture 2006. FAO Fisheries and Aquaculture Department, Rome, 2007.
- 1824 Food and Agriculture Organisation of the United Nations (FAO). 2007. The world's mangroves 1980-2005: a
1825 thematic study prepared in the framework of the Global Forest Resources Assessment 2005. Rome, 2007.
- 1826 Fourqurean, J. W., Marba, N., Duarte, C. M., Diaz-Almela, E., & Ruiz-Halpern, S. (2007). Spatial and temporal
1827 variation in the elemental and stable isotopic content of the seagrasses *Posidonia oceanica* and *Cymodocea*
1828 *nodosa* from the Illes Balears, Spain. *Marine Biology*, 151(1), 219-232.
- 1829 Fourqurean, J. W., Moore, T. O., Fry, B., & Hollibaugh, J. T. (1997). Spatial and temporal variation in C: N: P
1830 ratios, $\delta^{15}\text{N}$, and $\delta^{13}\text{C}$ of eelgrass *Zostera marina* as indicators of ecosystem processes, Tomales Bay,
1831 California, USA. *Marine Ecology Progress Series*, 157, 147-157.
- 1832 Fourqurean, J. W., Duarte, C. M., Kennedy, H., Marbà, N., Holmer, M., Mateo, M. A., & Serrano, O. (2012).
1833 Seagrass ecosystems as a globally significant carbon stock. *Nature Geoscience*, 5(7), 505-509.
- 1834 Fromard, H., Puig, E., Mougin, G., Marty, J., Bettoule, L., Cadamuro, F. (1998). Structure, above-ground
1835 biomass and dynamics of mangrove ecosystems: new data from French Guiana. *Oecologia* (115:), 39-43.
- 1836 Fujimoto, K., Imaya, A., Tabuchi, R., Kuramoto, S., Utsugi, H. 1999. Below-ground carbon storage of
1837 Micronesian mangrove forests. *Ecological Research* 14:409-413.
- 1838 Gallagher, J.L., Plumley, F.G. (1979) Underground biomass profiles and productivity in Atlantic coastal marshes.
1839 *American Journal of Botany* 66, 156-161.
- 1840 Gedan, K.B., Silliman, B. and Bertness, M.D. 2009. Centuries of Human-Driven Change in Salt Marsh
1841 Ecosystems. *Annual Review of Marine Science* 1:117–41.
- 1842 Gong, W. K., & Ong, J. E. 1990. Plant biomass and nutrient flux in a managed mangrove forest in Malaysia.
1843 *Estuarine, Coastal and Shelf Science*, 31(5), 519-530.
- 1844 Golley, F., Odum, H., Wilson, R. (1962). The structure and metabolism of a Puerto Rican red mangrove forest in
1845 May. *Ecological Society of America*, 43, 9-19.
- 1846 Golley, F. B., McGinnis, J. T., Clements, R. G., Child, G. I., & Duever, M. J. 1975. Mineral cycling in a tropical
1847 moist forest ecosystem.
- 1848 Goshima, S., & Peterson, C. H. (2012). Both below-and aboveground shoalgrass structure influence whelk
1849 predation on hard clams. *Marine Ecology Progress Series*, 451, 75.

Second Order Draft

- 1850 Green, E. P. and Short, F. T. 2003. World Atlas of Seagrasses. Prepared by the UNEP World Conservation
1851 Monitoring Centre, University of California Press, Berkeley, CA, USA.
- 1852 Groenendijk, A.M. (1984) Primary production of four dominant salt-marsh angiosperms in the SW Netherlands.
1853 Vegetatio 57, 143-152.
- 1854 Groenendijk, A.M., Vink-Lievaart, M.A. (1987) Primary production and biomass on a Dutch salt marsh:
1855 emphasis on the below-ground component. Vegetatio 70, 21-27.
- 1856 Gross, M.F., Hardisky, M.A., Wolf, P.L., Klemas, V. (1991) Relationship between aboveground and
1857 belowground biomass of *Spartina alterniflora* (smooth cordgrass) Estuaries 14, 180-191.
- 1858 Gross, M.F., Klemas, V., Levasseur, J.E. (1986) Biomass and structure of a *Spartina alterniflora* Loisel.-
1859 dominated salt marsh in France. Bulletin of the Torrey Botanical Society 113, 125-130.
- 1860 Guidetti, P., & Bussotti, S. (2002). Effects of seagrass canopy removal on fish in shallow Mediterranean seagrass
1861 (*Cymodocea nodosa* and *Zostera noltii*) meadows: a local-scale approach. Marine Biology, 140(3), 445-453.
- 1862 Gullström, M., Baden, S., & Lindegarth, M. (2012). Spatial patterns and environmental correlates in leaf-
1863 associated epifaunal assemblages of temperate seagrass (*Zostera marina*) meadows. Marine biology, 1-13.
- 1864 Hackney, J. W. (2003). Morphometric variability and allometric relationships in the seagrass *Thalassia*
1865 *testudinum* in Florida Bay (Doctoral dissertation, University of North Carolina).
- 1866 Halun, Z., Terrados, J., Borum, J., Kamp-Nielsen, L., Duarte, C. M., & Fortes, M. D. (2002). Experimental
1867 evaluation of the effects of siltation-derived changes in sediment conditions on the Philippine seagrass
1868 *Cymodocea rotundata*. Journal of Experimental Marine Biology and Ecology, 279(1), 73-87.
- 1869 Harris R.J, Mildbrandt E.C., Everham E.M. and Bovard B.D. 2010. The effects of reduced tidal flushing on
1870 mangrove structure and function across a disturbance gradient. Estuaries and Coasts 33:1176-1185.
- 1871 Hatala, J. A., Detto M., Sonnentag, O., Deverel, S. J., Verfaillie, J., Baldocchi, D. D. 2012. Greenhouse gas (CO₂,
1872 CH₄, H₂O) fluxes from drained and flooded agricultural peatlands in the Sacramento-San Joaquin Delta.
1873 Agriculture, Ecosystems and Environment 150: 1-18.
- 1874 Hatton, R. S., Patrick, W. H., & DeLaune, R. D. (1981). Sedimentation, nutrient accumulation, and early
1875 diagenesis in Louisiana Barataria Basin coastal marshes. In *Estuarine Comparisons 1982. Proceedings of the*
1876 *Sixth Biennial International Estuarine Research Conference*, Gleneden Beach, Oregon, November 1-6, 1981.
1877 p 255-267.
- 1878 Hebert, A. B., Morse, J. W., & Eldridge, P. M. (2007). Small-scale heterogeneity in the geochemistry of seagrass
1879 vegetated and non-vegetated estuarine sediments: causes and consequences. Aquatic Geochemistry, 13(1),
1880 19-39.
- 1881 Heck, K. L., & Thoman, T. A. (1984). The nursery role of seagrass meadows in the upper and lower reaches of
1882 the Chesapeake Bay. Estuaries and Coasts, 7(1), 70-92.
- 1883 Heijs, F. M. (1984). Annual biomass and production of epiphytes in three monospecific seagrass communities of
1884 *Thalassia hemprichii* (Ehrenb.) Aschers. Aquatic botany, 20(3), 195-218.
- 1885 Heilskov, A. C., Alperin, M. & Holmer, M. (2008). Benthic fauna bio-irrigation effects on nutrient regeneration
1886 in fish farm sediments. Journal of Experimental Marine Biology and Ecology, 339: 204–225.
- 1887 Heilskov, A. C. & Holmer, M. (2001) Effects of benthic fauna on organic matter mineralization in fish-farm
1888 sediments: importance of size and abundance. ICES Journal of Marine Science, 58: 427–434.
- 1889 Herbert, D. A. (1986). The growth dynamics of *Halophila hawaiiiana*. Aquatic botany, 23(4), 351-360.
- 1890 Herbert, D. A., & Fourqurean, J. W. (2008). Ecosystem structure and function still altered two decades after
1891 short-term fertilization of a seagrass meadow. Ecosystems, 11(5), 688-700.
- 1892 Herbert, D. A., & Fourqurean, J. W. (2009). Phosphorus availability and salinity control productivity and
1893 demography of the seagrass *Thalassia testudinum* in Florida Bay. Estuaries and coasts, 32(1), 188-201.
- 1894 Hillman, K., McComb, A. J., & Walker, D. I. (1995). The distribution, biomass and primary production of the
1895 seagrass *Halophila ovalis* in the Swan/Canning Estuary, Western Australia. Aquatic Botany, 51(1), 1-54.

- 1896 Holmer, M. & Kendrick, G. A. 2012. High Sulfide Intrusion in Five Temperate Seagrasses
1897 Growing Under Contrasting Sediment Conditions. *Estuaries and Coasts* 36, 1. 116-126.
- 1898 Holmer, M., Andersen, F. O., Nielsen, S. L. & Boschker, H. T. S. 2001. The importance of mineralization based
1899 on sulfate reduction for nutrient regeneration in tropical seagrass sediments. *Aquatic Botany* 71:1-17.
- 1900 Holmer, M., Duarte, C. M., Heilskov, A., Olesen, B., & Terrados, J. 2003. Biogeochemical conditions in
1901 sediments enriched by organic matter from net-pen fish farms in the Bolinao area, Philippines. *Marine*
1902 *pollution bulletin*, 46(11), 1470-1479.
- 1903 Holmer, M., Duarte, C. M., Boschker, H. T. S., & Barrón, C. 2004. Carbon cycling and bacterial carbon sources
1904 in pristine and impacted Mediterranean seagrass sediments. *Aquatic microbial ecology*, 36(3), 227-237.
- 1905 Holmer, M., Duarte, C. M., Heilskov, A., Olesen, B., & Terrados, J. (2002). Biogeochemical conditions in
1906 sediments enriched by organic matter from net-pen fish farms in the Bolinao area, Philippines. *Marine*
1907 *pollution bulletin*, 46(11), 1470-1479.
- 1908 Holmer, M., & Heilskov, A. C. 2008. Distribution and bioturbation effects of the tropical alpheid shrimp
1909 (*Alpheus macellarius*) in sediments impacted by milkfish farming. *Estuarine, Coastal and Shelf Science*,
1910 76(3), 657-667.
- 1911 Holmer, M., Wildish, D. & Hargrave, B. (2005). Organic Enrichment from Marine Finfish Aquaculture
1912 and Effects on Sediment Biogeochemical Processes. *Hdb Env Chem Vol. 5, Part M*: 181-206.
- 1913 Hopkinson, C.S., Gosselink, J.G., Parrando, R.T. (1978) Aboveground production of seven marsh plant species
1914 in coastal Louisiana. *Ecology* 59, 760-769.
- 1915 Hoque, A. T. M. R., Sharma, S., Suwa, R., Mori, S., & Hagihara, A. 2010. Seasonal variation in the size-
1916 dependent respiration of mangroves *Kandelia obovata*. *Mar. Ecol. Prog. Ser.*, 404, 31- 37.
- 1917 Howe, A. J., Rodriguez, J. F., Saco, P. M. 2009. Surface evolution and carbon sequestration in disturbed and
1918 undisturbed wetland soils of the Hunter estuary, southeast Australia. *Estuarine, Coastal and Shelf Science* 84:
1919 75-83.
- 1920 Hu, Z., Lee, JW, Chandran, K, Kim, S, and Khanal, K. 2012. Nitrous oxide (N₂O) emission from aquaculture: a
1921 review. *Environmental Science and Technology* 46:6470-6480.
- 1922 Hussein, A. H., M. C. Rabenhorst, et al. (2004). Modeling of carbon sequestration in coastal marsh soils. *Soil*
1923 *Science Society of America Journal* 68: 1786-1795.
- 1924 Hussey, A., Long, S.P. (1982) Seasonal changes in weight of above- and below-ground vegetation and dead
1925 plant material in a salt marsh at Colne Point, Essex. *Journal of Ecology* 70, 757-771.
- 1926 IPCC 2003, Good Practice Guidance for Land Use, Land-Use Change and Forestry. Penman J., Gytarsky M.,
1927 Hiraishi T., Krug T., Kruger D., Pipatti R., Buendia L., Miwa K., Ngara T., Tanabe K., Wagner F. (eds.).
1928 Intergovernmental Panel on Climate Change (IPCC), IPCC/IGES, Hayama, Japan.
- 1929 IPCC 2006. 2006 IPCC Guidelines for National Greenhouse Gas Inventories, Prepared by the National
1930 Greenhouse Gas Inventories Programme, Eggleston H.S., Buendia L., Miwa K., Ngara T. and Tanabe K.
1931 (eds.). Intergovernmental Panel on Climate Change (IPCC), IPCC/IGES, Hayama, Japan.
- 1932 Ismail, N. (1993). Preliminary study of the seagrass flora of Sabah, Malaysia. *Pertanika Journal of Tropical*
1933 *Agricultural Science*, 16(2), 111-118.
- 1934 Iverson, R. L., & Bittaker, H. F. (1986). Seagrass distribution and abundance in eastern Gulf of Mexico coastal
1935 waters. *Estuarine, Coastal and Shelf Science*, 22(5), 577-602.
- 1936 Jacobs, R. P. W. M. (1979). Distribution and aspects of the production and biomass of eelgrass, *Zostera marina*
1937 L., at Roscoff, France. *Aquatic Botany*, 7, 151-172.
- 1938 James, N. P., Bone, Y., Brown, K. M., & Cheshire, A. (2009). Calcareous epiphyte production in cool-water
1939 carbonate seagrass depositional environments-southern Australia. *International Association of*
1940 *Sedimentology Special Publication*, 41, 123-148.
- 1941 Jeffrey, D.W., Hayes, M., (2005) Net primary productivity of intertidal systems: the Dublin Bay example, in:
1942 Wilson, J.G. (Ed.), *The Intertidal Ecosystem: The Value of Ireland's Shores*. Royal Irish Academy, Dublin,
1943 pp. 45-57.

Second Order Draft

- 1944 Jeffries, R.L., Davy, A.J., Rudmik, T. (1981) Population biology of the salt marsh annual *Salicornia europaea*
1945 agg. *Journal of Ecology* 69, 17-31.
- 1946 Jensen, S., & Bell, S. (2001). Seagrass growth and patch dynamics: cross-scale morphological plasticity. *Plant*
1947 *Ecology*, 155(2), 201-217.
- 1948 Jones, D. A.; Nithyanandan, M.; Williams, I 2012 Sabah Al-Ahmad Sea City Kuwait: development of a
1949 sustainable man-made coastal ecosystem in a saline desert. *AQUATIC ECOSYSTEM HEALTH &*
1950 *MANAGEMENT* Volume: 15 Special Issue: SI Supplement: 1 Pages: 84-92
- 1951 Juliana, W.A. and Nizam, M.S. 2004. Forest structure and above-ground biomass of two mangrove forest
1952 communities in Matang. *In:* (M.I. Shaharuddin, A. Muda, R. Ujang, A.B. Kamaruzaman, K.L. Lim, S. Rosli,
1953 J.M. Som and A. Latiff, eds) *Proceedings of the National Conference on Sustainable Management of the*
1954 *Matang Mangroves-100 Years and Beyond*. Ipoh, Perak, Malaysia. pp. 235-244.
- 1955 Kairo, J. G., Lang'at, J., Dahdouh-Guebas, F., Bosire, J., Karachi, M. (2008). Structural development and
1956 productivity of replanted mangrove plantations in Kenya. *Forest Ecology and Management*(255), 2670-2677.
- 1957 Kaldy, J. E. (2006). Production ecology of the non-indigenous seagrass, dwarf eelgrass (*Zostera japonica* Ascher.
1958 & Graeb.), in a Pacific Northwest estuary, USA. *Hydrobiologia*, 553(1), 201-217.
- 1959 Kaldy, J. E., & Dunton, K. H. (2000). Above-and below-ground production, biomass and reproductive ecology
1960 of *Thalassia testudinum* (turtle grass) in a subtropical coastal lagoon. *Marine ecology. Progress series*, 193,
1961 271-283.
- 1962 Kauffman, J. B. Heider, C., Cole, T. G. (2011) Ecosystem carbon stocks of Micronesian mangrove forests.
1963 *Wetlands*, 31: 343-352.
- 1964 Kearney, M. S., & Stevenson, J. C. (1991). Island land loss and marsh vertical accretion rate evidence for
1965 historical sea-level changes in Chesapeake Bay. *Journal of Coastal Research*, 403-415.
- 1966 Kentula, M. E., & David McIntire, C. (1986). The autecology and production dynamics of eelgrass (*Zostera*
1967 *marina* L.) in Netarts Bay, Oregon. *Estuaries and Coasts*, 9(3), 188-199.
- 1968 Kenworthy, J. W., & Thayer, G. W. (1984). Production and decomposition of the roots and rhizomes of
1969 seagrasses, *Zostera marina* and *Thalassia testudinum*, in temperate and subtropical marine ecosystems.
1970 *Bulletin of Marine Science*, 35(3), 364-379.
- 1971 Kerr, E. A., & Strother, S. (1989). Seasonal changes in leaf growth rate of *Zostera muelleri* Irmisch ex Aschers.
1972 In South-Eastern Australia. *Aquatic botany*, 33(1), 131-140.
- 1973 Khan, M. N. I., Suwa, R., & Hagihara, A. 2009. Biomass and aboveground net primary production in a
1974 subtropical mangrove stand of *Kandelia obovata* (S., L.) Yong at Manko Wetland, Okinawa, Japan. *Wetlands*
1975 *Ecology and Management*, 17(6), 585-599.
- 1976 Kim, S. H., Kim, Y. K., Park, S. R., Li, W. T., Lee, K. S. 2012. Growth dynamics of the seagrass *Halophila*
1977 *nipponica*, recently discovered in temperate coastal waters of the Korean peninsula. *Mar Biol*, 159:255–267.
- 1978 Kirby, C.J., Gosselink, J.G. (1976) Production in a Louisiana Gulf Coast *Spartina alterniflora* marsh. *Ecology*
1979 57, 1052-1059.
- 1980 Kirkman, H., & Cook, I. H. (1987). Distribution and leaf growth of *Thalassodendron pachyrhizum* den Hartog in
1981 Southern Western Australia. *Aquatic botany*, 27(3), 257-266.
- 1982 Kirkman, H., & Reid, D. D. (1979). A study of the role of the seagrass *Posidonia australis* in the carbon budget
1983 of an estuary. *Aquatic Botany*, 7, 173-183.
- 1984 Kirkman, H., Cook, I. H., & Reid, D. D. (1982). Biomass and growth of *Zostera capricorni* aschers. in port
1985 hacking, NSW, Australia. *Aquatic Botany*, 12, 57-67.
- 1986 Kirui, B., Kairo, J. G., Karachi, M. (2006.). Allometric Equations for Estimating Above Ground Biomass of
1987 *Rhizophora mucronata* Lamk. (Rhizophoraceae) Mangroves at Gazi Bay, Kenya. *Western Indian Ocean J.*
1988 *Mar. Sci.*, 5(1): , 27-34
- 1989 Kirwan, M. L., Murray, A.B., Donnelly, J.B., Corbett, D.R. (2011) Rapid wetland expansion during European
1990 settlement and its implication for marsh survival under modern sediment delivery rates. *Geology* 39: 507-510.

- 1991 Kistritz, R.U., Hall, K.J., Yesaki, I. (1983) Productivity, detritus flux, and nutrient cycling in a *Carex lyngbyei*
1992 tidal marsh. *Estuaries* 6, 227-236.
- 1993 Koch, B. P., Souza Filho, P. W., Behling, H., Cohen, M. C., Kattner, G., Rullkötter, J., Scholz-Böttcher B.M. &
1994 Lara, R. J. 2011. Triterpenols in mangrove sediments as a proxy for organic matter derived from the red
1995 mangrove (*Rhizophora mangle*). *Organic Geochemistry*, 42(1), 62-73.
- 1996 Komiyama, A., Moriya, H., Prawiroatmodjo, S., Toma, T., & Ogino, K. (1988). Primary productivity of
1997 mangrove forest. *Biological system of mangroves*. Ehime University, Ehime, Japan, 96-97.
- 1998 Komiyama, A., Havanond, S., Siraswatt, W., Muchida, Y., Fujimoto, K. (2000). Top/root biomass ratio of a
1999 secondary mangrove (*Ceriops tagal* (Perr.) C. B Rob.) forest. *Forest Ecology and Management*(139), 127-
2000 134.
- 2001 Komiyama, A., Ogino, K., Aksornkoae, S., Sabhasri, S. (1987). Root biomass of a mangrove forest in southern
2002 Thailand. 1. Estimation by the trench method and the zonal structure of root biomass. *Journal of Tropical*
2003 *Ecology*(3), 97-108.
- 2004 Komiyama, A., Ong, J., Pongporm, S. (2008). Allometry, biomass, and productivity of mangrove forests: A
2005 review. *Aquatic Botany*(89), 128-137.
- 2006 Kowalski, J. L., DeYoe, H. R., & Allison, T. C. (2009). Seasonal production and biomass of the seagrass,
2007 *Halodule wrightii* Aschers.(shoal grass), in a subtropical Texas lagoon. *Estuaries and coasts*, 32(3), 467-482.
- 2008 Kraemer, G. P., and Alberte, R. S. 1995. Impact of daily photosynthetic period on protein synthesis and
2009 carbohydrate stores in *Zostera marina* L. (eelgrass) roots: implications for survival in light-limited
2010 environments. *Journal of Experimental Biology and Ecology* 185: 191-202.
- 2011 Krauss KW, TW Doyle, RR Twilley, TJ Smith, KT Whelan, JK Sullivan. 2005. Woody debris in the mangrove
2012 forests of south Florida. *Biotropica* 37: 9-15
- 2013 Kreuzwieser J, Buchholz J, and Rennenberg H. 2003. Emission of methane and nitrous oxide by Australian
2014 Krithika, mangrove ecosystems. *Plant Biology* 5: 423–431.
- 2015 Kristensen, E., S. Bouillon, T Dittmar, C Marchand. 2008. Organic carbon dynamics in mangrove ecosystems: a
2016 review. *Aquatic Botany* 89: 201-219.
- 2017 Kristensen, E., Flindt, M.R., Ulomi, S., Borges, A.V., Abril, G., Bouillon S. (2008) Emission of CO₂ and CH₄ to
2018 the atmosphere by sediments and open waters in two Tanzanian mangrove forests. *Marine Ecology Progress*
2019 *Series*, 370, 53-67.
- 2020 Krithika, K, Purvaja, R, Ramesh, R. 2008. Fluxes of methane and nitrous oxide from an Indian mangrove.
2021 *Current Science* 94: 218-224.
- 2022 Kungvankij, P., Pudadera Jr, B. J., Tiro Jr, L. B., Potestas, I. O., & Chua, T. E. 1986. An improved traditional
2023 shrimp culture technique for increasing pond yield.
- 2024 Kurz, W. A., Apps, M., Banfield, E., & Stinson, G. 2002. Forest carbon accounting at the operational scale. *The*
2025 *Forestry Chronicle*, 78(5), 672-679.
- 2026 Kusmana, C., Sabiham, S., Abe, K., & Watanabe, H. (1992). An estimation of above ground tree biomass of a
2027 mangrove forest in East Sumatra, Indonesia. *Tropics*, 1(4), 243-257.
- 2028 Larkum, A. W. D., Collett, L. C., & Williams, R. J. (1984). The standing stock, growth and shoot production of
2029 *Zostera capricorni* aschers. in Botany Bay, New South Wales, Australia. *Aquatic botany*, 19(3), 307-327.
- 2030 Larkum, A.W.D. and West R.J. 1990 Long-term changes of seagrass meadows in Botany Bay, Australia.
2031 *Aquatic Botany* 37: 55-70.
- 2032 Larned, S. T. (2003). Effects of the invasive, nonindigenous seagrass *Zostera japonica* on nutrient fluxes
2033 between the water column and benthos in a NE Pacific estuary. *Marine Ecology Progress Series*, 254, 69-80.
- 2034 Latimer, J.S.; Rego, S.A.2010. Empirical relationship between eelgrass extent and predicted watershed-derived
2035 nitrogen loading for shallow New England estuaries. *Estuarine, Coastal and Shelf Science* 90, 231-240.

Second Order Draft

- 2036 Laugier, T., Rigollet, V., & de Casabianca, M. L. (1999). Seasonal dynamics in mixed eelgrass beds, *Zostera*
 2037 *marina* L. and *Z. noltii* Hornem., in a Mediterranean coastal lagoon (Thau lagoon, France). *Aquatic botany*,
 2038 63(1), 51-69.
- 2039 Lebreton, B., Richard, P., Radenac, G., Bordes, M., Br  ret, M., Arnaud, C. & Blanchard, G. F. (2009). Are
 2040 epiphytes a significant component of intertidal *Zostera noltii* beds?. *Aquatic Botany*, 91(2), 82-90.
- 2041 Lee, K. S., & Dunton, K. H. (1996). Production and carbon reserve dynamics of the seagrass *Thalassia*
 2042 *testudinum* in Corpus Christi Bay, Texas, USA. *Marine ecology progress series*. Oldendorf, 143(1), 201-210.
- 2043 Lee, K. S., Park, S. R., & Kim, J. B. (2005). Production dynamics of the eelgrass, *Zostera marina* in two bay
 2044 systems on the south coast of the Korean peninsula. *Marine Biology*, 147(5), 1091-1108.
- 2045 Lee, S. Y. (1997). Annual cycle of biomass of a threatened population of the intertidal seagrass *Zostera japonica*
 2046 in Hong Kong. *Marine Biology*, 129(1), 183-193.
- 2047 Lee, S. Y., Oh, J. H., Choi, C. I., Suh, Y., & Mukai, H. (2005b). Leaf growth and population dynamics of
 2048 intertidal *Zostera japonica* on the western coast of Korea. *Aquatic botany*, 83(4), 263-280.
- 2049 Lee, S. Y. 1990. Primary productivity and particulate organic matter flow in an estuarine mangrove-wetland in
 2050 Hong Kong. *Marine Biology*, 106(3), 453-463.
- 2051 Leendertse P.C., Rozema J. and Andrea A. 1997. Effects of nitrogen addition on the growth of the salt marsh
 2052 grass *Elymus athericus*. *Estuarine, Coastal and Shelf Science* 3: 35-40.
- 2053 Lewis, F. G. (1987). Crustacean epifauna of seagrass and macroalgae in Apalachee Bay, Florida, USA. *Marine*
 2054 *Biology*, 94(2), 219-229.
- 2055 Liao B.W., Zheng D.Z. and Zheng S.F. 1990. Studies on the Biomass of *Sonneratia caseolaris* stand. *Forest*
 2056 *Research* 1: 47-54(in Chinese with English abstract).
- 2057 Liikanen, A, Silvennoinen, H. Karvo, A, Rantakokko, P, Martikainen, PJ. 2009. *Boreal Environment Research*
 2058 14: 351-368.
- 2059 Lilleb  , A. I., Flindt, M. R., Pardal, M. A., & Marques, J. C. (2006). The effect of *Zostera noltii*, *Spartina*
 2060 *maritima* and *Scirpus maritimus* on sediment pore-water profiles in a temperate intertidal estuary.
 2061 *Hydrobiologia*, 555(1), 175-183.
- 2062 Lin, P. 1989. Biomass and element cycle of *Kandelia* forest, China. *Mangrove Research Papers* (1980-1989),
 2063 Xiamen Univ. Press, Xiamen, 143-149.
- 2064 Lin P. and Lu C.Y. 1990. Biomass and productivity of *Bruguiera sexangula* mangrove forest in Hainan Island,
 2065 China. *Journal of Xiamen University (Natural Science)* 29: 209-213 (In Chinese with English abstract)
- 2066 Lindeboom, H. J., & Sandee, A. J. J. (1989). Production and consumption of tropical seagrass fields in eastern
 2067 Indonesia measured with bell jars and microelectrodes. *Netherlands Journal of Sea Research*, 23(2), 181-190.
- 2068 Linthurst, R.A., Reimold, R.J. (1978) Estimated net aerial primary production for selected estuarine angiosperms
 2069 in Maine, Delaware, and Georgia. *Ecology* 59, 945-955.
- 2070 Lipkin, Y. (1979). Quantitative aspects of seagrass communities, particularly of those dominated by *Halophila*
 2071 *stipulacea*, in Sinai (Northern Red Sea). *Aquatic Botany*, 7, 119-128.
- 2072 Livesley, S., Andrusiak, S., & Idczak, D. 2010. Soil greenhouse gas exchange and carbon stocks in natural and
 2073 managed ecosystems of the Mornington Peninsula. Final Report to the Mornington Peninsula Shire Council,
 2074 Melbourne, Australia.
- 2075 Livingstone, D.C., Patriquin, D.G. (1981) Belowground growth of *Spartina alterniflora* Loisel.: habitat,
 2076 functional biomass and non-structural carbohydrates. *Estuarine Coastal and Shelf Science* 12, 579-587.
- 2077 Loomis, M. J., & Craft, C. B. (2010). Carbon sequestration and nutrient (nitrogen, phosphorus) accumulation in
 2078 river-dominated tidal marshes, Georgia, USA. *Soil Science Society of America Journal*, 74(3), 1028-1036.
- 2079 Long, W. L., Mellors, J. E., & Coles, R. G. (1993). Seagrasses between Cape York and Hervey Bay, Queensland,
 2080 Australia. *Marine and Freshwater Research*, 44(1), 19-31.

- 2081 Longstaff, B. J., & Dennison, W. C. (1999). Seagrass survival during pulsed turbidity events: the effects of light
2082 deprivation on the seagrasses *Halodule pinifolia* and *Halophila ovalis*. *Aquatic Botany*, 65(1), 105-121.
- 2083 Longstaff, B. J., Loneragan, N. R., O'Donohue, M. J., & Dennison, W. C. (1999). Effects of light deprivation on
2084 the survival and recovery of the seagrass *Halophila ovalis* (R. Br.) Hook. *Journal of experimental marine*
2085 *biology and ecology*, 234(1), 1-27.
- 2086 Lovelock, CE, Ruess, RW, Feller, IC. 2011. CO₂ efflux from cleared mangrove peat. *PLOS One* 6: 1-4.
- 2087 Lu, C., Wong, Y., Tam, N.F.Y., Ye Y., Cui, S., Lin, P. (1998). Preliminary studies on methane fluxes in Hainan
2088 mangrove communities *Chinese Journal of Oceanology and Limnology*: 16, 64-71.
- 2089 Lugo, A., Snedaker, S. (1974). The Ecology of Mangroves. *Annual Review of Ecology and Systematics*, 5,: 39-
2090 64.
- 2091 Mackey, A. P. (1993). Biomass of the Mangrove *Avicennia marina* (Forsk.) Vierh. Near Brisbane, South-eastern
2092 Queensland. *Australian Journal of Marine and Freshwater Research* .(44:), 721-5.
- 2093 Madkour, F. F. and Gaballah, M. M. 2012. Phytoplankton assemblage of a solar saltern in Port Fouad, Egypt.
2094 *Oceanologia* 54: 687-700.
- 2095 Mahall, B.E., Park, R.B. (1976) The ecotone between *Spartina foliosa* Trin. and *Salicornia virginica* L. in salt
2096 marshes of Northern San Francisco Bay: II. Soil water and salinity. *Journal of Ecology* 64, 793-809.
- 2097 Mall, L. P., Singh, V. P., Garge, A. 1(1991). Study of biomass, litterfall, litter decomposition and soil respiration
2098 in monogeneric mangrove. *Tropical Ecology* .(32:), 144-152.
- 2099 Mann, K. H. (1972). Ecological energetics of the seaweed zone in a marine bay on the Atlantic coast of Canada.
2100 I. Zonation and biomass of seaweeds. *Marine biology*, 12(1), 1-10.
- 2101 Marbà, N., & Duarte, C. M. (2001). Growth and sediment space occupation by seagrass *Cymodocea nodosa*
2102 roots.
- 2103 Marba, N., and D.I. Walker. 1999. Growth, flowering, and population dynamics of temperate Western Australian
2104 seagrasses. *Marine Ecology Progress Series* 184: 105–118.
- 2105 Marchand, C., J.R. Disnar, E. Lallier-Vesges, N. Lottier. 2005. Early diagenesis of carbohydrates and lignin in
2106 mangrove sediments subject to variable redox conditions (French Guiana). *Geochim. Cosmochim. Acta*
2107 69:131-142.
- 2108 Marchand, C., E. Lallier-Verges, F. Baltzer. 2003. The composition of sedimentary organic matter in relation to
2109 the dynamic features of a mangrove-fringed coast in French Guiana. *ECSS* 56: 119-130
- 2110 Markewich, H. W., Wysocki, D. A., Pavich, M. J., Rutledge, E. M., Millard Jr, H. T., Rich, F. J., ... & McGeehin,
2111 J. P. (1998). Paleopedology plus TL,< sup> 10</sup> Be, and< sup> 14</sup> C dating as tools in
2112 stratigraphic and paleoclimatic investigations, Mississippi River Valley, USA. *Quaternary International*, 51,
2113 143-167.
- 2114 Martin, S., Clavier, J., Guarini, J. M., Chauvaud, L., Hily, C., Grall, J., & Richard, J. (2005). Comparison of
2115 *Zostera marina* and maerl community metabolism. *Aquatic botany*, 83(3), 161-174.
- 2116 Mascaro, O., Valdemarsen, T., Holmer, M., Perez, M., & Romero, J. (2009). Experimental manipulation of
2117 sediment organic content and water column aeration reduces *Zostera marina* (eelgrass) growth and survival.
2118 *Journal of Experimental Marine Biology and Ecology*, 373(1), 26-34.
- 2119 Mateo, M. A., & Romero, J. (1997). Detritus dynamics in the seagrass *Posidonia oceanica*: elements for an
2120 ecosystem carbon and nutrient budget. *Marine Ecology Progress Series*, 151(1), 43-53.
- 2121 Matsui, N. (1998). Estimated stocks of organic carbon in mangrove roots and sediments in
2122 Hinchinbrook Channel, Australia. *Mangroves and Salt Marshes* 2: 199–204.
- 2123 McCaffrey, R. J., & Thomson, J. (1980). A record of the accumulation of sediment and trace metals in a
2124 Connecticut salt marsh. *Advances in geophysics*, 22, 165-236.
- 2125 McDermid, K. J., & Edward, A. (1999). Seagrass community composition and biomass at Nahpali Island,
2126 Pohnpei. *MICRONESICA-AGANA*-, 31, 255-262.

Second Order Draft

- 2127 McGlathery, K.J, Reynolds, L.K, Cole, L.W, Orth, R.J, Marion, S.R, Schwarzschild, A. 2012. Recovery trajectories
2128 during state change from bare sediment to eelgrass dominance. *Marine Ecology Progress Series* 448: 209-221.
- 2129 McKenzie, L. J. (1994). Seasonal changes in biomass and shoot characteristics of a *Zostera capricorni* Aschers.
2130 dominant meadow in Cairns Harbour, northern Queensland. *Marine and Freshwater Research*, 45(7), 1337-
2131 1352.
- 2132 McMahan, C. A. (1968). Biomass and salinity tolerance of shoalgrass and manatee grass in Lower Laguna Madre,
2133 Texas. *The Journal of Wildlife Management*, 501-506.
- 2134 McMahon, K., & Lavery, P. (2008). The responses of *Amphibolis griffithii* to reduced light availability. Final
2135 report on the Strategic Research Fund for the Marine Environment (SRFME) Collaborative Research Project:
2136 Ecophysiology of benthic primary producers, Centre for Marine Ecosystems Research, Edith Cowan
2137 University, Joondalup, Western Australia.
- 2138 McRoy, C. P. (1974). Seagrass productivity: Carbon uptake experiments in eelgrass, *Zostera marina*.
2139 *Aquaculture*, 4, 131-137.
- 2140 Medeiros, T., Sampaio, E. (2008). Allometry of above-ground biomasses in mangrove species in Itamaraca,
2141 Pernambuco, Brazil. *Wetlands Ecology and Management*, 16(3), 323-330.
- 2142 Megonigal, J.P. and Neubauer, S.C. (2009). Chapter 19 Biogeochemistry of tidal freshwater wetlands. In:
2143 Perillo, G.M.E, Wolanski, E., Cahoon, D.R., Brinson, M.M. (editors) *Coastal Wetlands: An Integrated*
2144 *Ecosystem Approach*. Elsevier.
- 2145 Meling-López, A. E., & Ibarra-Obando, S. E. (1999). Annual life cycles of two *Zostera marina* L. populations in
2146 the Gulf of California: contrasts in seasonality and reproductive effort. *Aquatic Botany*, 65(1), 59-69.
- 2147 Mellors, J., Marsh, H., Carruthers, T. J., & Waycott, M. (2002). Testing the sediment-trapping paradigm of
2148 seagrass: Do seagrasses influence nutrient status and sediment structure in tropical intertidal environments?.
2149 *Bulletin of marine science*, 71(3), 1215-1226.
- 2150 Mendelssohn, I.A., Marcellus, K.L. (1976) Angiosperm production of three Virginia marshes in various salinity
2151 and soil nutrient regimes. *Chesapeake Science* 17, 15-23.
- 2152 Moncreiff, C. A., Sullivan, M. J., & Daehnick, A. E. (1992). Primary production dynamics in seagrass beds of
2153 Mississippi Sound: the contributions of seagrass, epiphytic algae, sand microflora, and phytoplankton.
2154 *Marine Ecology-Progress Series*, 87, 161-161.
- 2155 Morgan, M. D., & Kitting, C. L. (1984). Productivity and utilization of the seagrass *Halodule wrightii* and its
2156 attached epiphytes. *Limnology and Oceanography*, 1066-1076.
- 2157 Moriarty, D. J. W., Roberts, D. J., & Pollard, P. C. (1990). Primary and bacterial productivity of tropical seagrass
2158 communities in the Gulf of Carpentaria, Australia. *Marine Ecology Progress Series* MESED, 61(1/2).
- 2159 Morris, J. T., Edwards, J., Crooks, S., Reyes, E. 2012. Assessment of Carbon Sequestration Potential in Coastal
2160 Wetlands. In *Recarbonization of the Biosphere*, Eds. Lal, R., Lorenz, K., Hüttel, R. F., Schneider, B. U., von
2161 Braun, J., Springer, Dordrecht, 559p.
- 2162 Morris, J.T., Haskin, B. (1990) A 5-yr record of aerial primary production and stand characteristics of *Spartina*
2163 *alterniflora*. *Ecology* 71, 2209-2217.
- 2164 Morris, J. T., & Jensen, A. (2003). The carbon balance of grazed and non-grazed *Spartina anglica* saltmarshes at
2165 Skallingen, Denmark. *Journal of Ecology*, 86(2), 229-242.
- 2166 Moseman-Valtierra S., Gonzalez R., Kroeger K., Tang J., Chun W., Crusius J., Bratton J., Green A. and
2167 Shelton J. 2011. Short-term nitrogen additions can shift a coastal wetland from a sink to a source of
2168 N₂O. *Atmospheric Environment* 45: 4390-4397.
- 2169 Mukai, H., Aioi, K., Koike, I., Iizumi, H., Ohtsu, M., & Hattori, A. (1979). Growth and organic production of
2170 eelgrass (*Zostera marina* L.) in temperate waters of the Pacific coast of Japan. I. Growth analysis in spring-
2171 summer. *Aquatic Botany*, 7, 47-56.
- 2172 Murray, L., & Wetzel, R. L. (1987). Oxygen production and consumption associated with the major autotrophic
2173 components in two temperate seagrass communities. *Mar Ecol Prog Ser*, 38, 231-239.

- 2174 Neves, J.P., Ferreira, L.F., Simoes, M.P., Gazarini, L.C. (2007) Primary production and nutrient content in two
2175 salt marsh species, *Atriplex portulacoides* L. and *Limoniastrum monopetalum* L., in Southern Portugal.
2176 *Estuaries and Coasts* 30, 459-468.
- 2177 Nienhuis, P. H., Coosen, J., & Kiswara, W. (1989). Community structure and biomass distribution of seagrasses
2178 and macrofauna in the Flores Sea, Indonesia. *Netherlands Journal of Sea Research*, 23(2), 197-214.
- 2179 Nixon, S. W., & Oviatt, C. A. (1972). Preliminary measurements of midsummer metabolism in beds of eelgrass,
2180 *Zostera marina*. *Ecology*, 150-153.
- 2181 Norhayati, A. and A. Latiff, 2001. Biomass and species composition of a mangrove forest in Pulau Langkawi,
2182 Malaysia. *Malaysia Applied Biol.*, 30: 75-80.
- 2183 Odum, H. T. (1963). Productivity measurements in Texas turtle grass and the effects of dredging an intracoastal
2184 channel. *Publications of the Institute of Marine Science*, 9, 48-58.
- 2185 Oenema, O., & DeLaune, R. D. (1988). Accretion rates in salt marshes in the Eastern Scheldt, south-west
2186 Netherlands. *Estuarine, Coastal and Shelf Science*, 26(4), 379-394.
- 2187 Ogden, J. C., & Ogden, N. B. (1982). A preliminary study of two representative seagrass communities in Palau,
2188 Western Caroline Islands (Micronesia). *Aquatic Botany*, 12, 229-244.
- 2189 Olesen, B. (1999). Reproduction in Danish eelgrass (*Zostera marina* L.) stands: size-dependence and biomass
2190 partitioning. *Aquatic botany*, 65(1), 209-219.
- 2191 Olesen, B., & Sand-Jensen, K. (1994). Biomass-density patterns in the temperate seagrass *Zostera marina*.
2192 *Marine Ecology-Progress Series*, 109, 283-283.
- 2193 Oleson, B., & Sand-Jensen, K. (1993). Seasonal acclimitization of eelgrass *Zostera marina* growth to light.
2194 *Marine Ecology Progress Series MESED*., 94(1).
- 2195 Ong, E. 1982. Mangroves and aquaculture in Malaysia. *Ambio*, 11(5), 252-257.
- 2196 Oren A 2002 Molecular ecology of extremely halophilic Archaea and Bacteria.FEMS MICROBIOLOGY
2197 ECOLOGY Volume: 39 Issue: 1 Pages: 1-7
- 2198 Oren A 2009 Saltern evaporation ponds as model systems for the study of primary production processes under
2199 hypersaline conditions.Aquat Microb Ecol Volume: 56 Issue: 2-3 Pages: 193-204
- 2200 Orson, R. A., Warren, R. S., & Niering, W. A. (1998). Interpreting sea level rise and rates of vertical marsh
2201 accretion in a southern New England tidal salt marsh. *Estuarine, Coastal and Shelf Science*, 47(4), 419-429.
- 2202 Orth, R. J., & Moore, K. A. (1986). Seasonal and year-to-year variations in the growth of *Zostera marina* L.
2203 (eelgrass) in the lower Chesapeake Bay. *Aquatic Botany*, 24(4), 335-341.
- 2204 Orth, R. J.; Moore, .K. A.; Marion, S. R., Wilcox, D. J., Parrish, D. B. 2012 Seed addition facilitates eelgrass
2205 recovery in a coastal bay system . *Marine Ecology Progress Series* Volume: 448 Pages: 177-195
- 2206 Ortiz-Milan SM: "Project of recovery the biological conditions of the production system in saltworks of
2207 Industria Salinera de Yucatan S.A. de C.V. (ISYSA) damaged by the hurricane isidore in September of 2002",
2208 In *CEISSA2006*: (Oct 2006)
- 2209 Osborne, T. Z., Bruland, G. L., Newman, S., Reddy, K. R., & Grunwald, S. (2011). Spatial distributions and eco-
2210 partitioning of soil biogeochemical properties in the Everglades National Park. *Environmental monitoring*
2211 *and assessment*, 183(1), 395-408.
- 2212 Osland M.J., Spivak A.C., Nestlerode J.A., Lessmann J.M., Almario A.E., Heitmuller P.T., Russell M.J., Krauss
2213 K.W., Alvarez F., Dantin D.D., Harvey J.E., From A.S, Cormier N., Stagg C.L. 2012. Ecosystem
2214 development after mangrove wetland creation: plant-soil change across a 20-year chronosequence.
2215 *Ecosystems* DOI: 10.1007/s10021-012-9551-1
- 2216 Ott, J. A. (2008). Growth and Production in *Posidonia Oceanica* (L.) Delile*. *Marine Ecology*, 1(1), 47-64.
- 2217 Pa'Ez-osun F 2001AThe Environmental Impact of Shrimp Aquaculture: Causes, Effects, and Mitigating
2218 Alternatives. *Environmental Management* Vol. 28, No. 1, pp. 131-140
- 2219 Page, K. L. & Dal, R. C. 2011. Contribution of natural and drained wetland systems to carbon stocks, CO₂, N₂O,
2220 and CH₄ fluxes: an Australian perspective. *Soil Research* 49: 377-388.

Second Order Draft

- 2221 Paling, E. I., & McComb, A. J. (2000). Autumn biomass, below-ground productivity, rhizome growth at bed
2222 edge and nitrogen content in seagrasses from Western Australia. *Aquatic Botany*, 67(3), 207-219.
- 2223 Park, S. R., Kim, Y. K., Kim, J. H., Kang, C. K., & Lee, K. S. (2011). Rapid recovery of the intertidal seagrass
2224 *Zostera japonica* following intense Manila clam (*Ruditapes philippinarum*) harvesting activity in
2225 Korea. *Journal of Experimental Marine Biology and Ecology*, 407(2), 275-283.
- 2226 Patrick Jr, W. H., & DeLaune, R. D. (1990). Subsidence, accretion, and sea level rise in south San Francisco Bay
2227 marshes. *Limnology and Oceanography*, 1389-1395.
- 2228 Paynter, C. K., Cortés, J., & Engels, M. (2001). Biomass, productivity and density of the seagrass *Thalassia*
2229 *testudinum* at three sites in Cahuita National Park, Costa Rica. *Rev. Biol. Trop.*, 49(Suppl 2), 265-272.
- 2230 Pedersen, M. F., & Borum, J. (1993). An annual nitrogen budget for a seagrass *Zostera marina* population.
2231 *Marine Ecology-Progress Series*, (101), 169-177.
- 2232 Pennings S.C., Stanton L.E. and Stephen Brewer J. 2002. Nutrient effects on the composition of salt marsh plant
2233 communities along the southern Atlantic and Gulf coasts of the United States. *Estuaries* 25: 1164–1173.
- 2234 Pérez-Lloréns, J. L., & Niell, F. X. (1993). Seasonal dynamics of biomass and nutrient content in the intertidal
2235 seagrass *Zostera noltii* Hornem. from Palmones River estuary, Spain. *Aquatic botany*, 46(1), 49-66.
- 2236 Pérez-Ruzafa, A., Marcos, C., Bernal, C. M., Quintino, V., Freitas, R., Rodrigues, A. M., & Pérez-Ruzafa, I. M.
2237 (2012). *Cymodocea nodosa* vs. *Caulerpa prolifera*: causes and consequences of a long term history of
2238 interaction in macrophyte meadows in the Mar Menor coastal lagoon (Spain, Southwestern Mediterranean).
2239 *Estuarine, Coastal and Shelf Science*.
- 2240 Pérez, M., & Romero, J. (1994). Growth Dynamics, Production, and Nutrient Status of the Seagrass *Cymodocea*
2241 *nodosa* in a Mediterranean Semi-Estuarine Environment. *Marine Ecology*, 15(1), 51-64.
- 2242 Pergent, G., Romero, J., Pergent-Martini, C., Mateo, M. A., & Boudouresque, C. F. (1994). Primary production,
2243 stocks and fluxes in the Mediterranean seagrass *Posidonia oceanica*. *Marine Ecology-Progress Series*, 106,
2244 139-139.
- 2245 Perillo G.M.E., Wolanski E., Cahoon D.R., Brinson M.M. 2009. Coastal Wetlands: An Integrated Ecosystem
2246 Approach. Elsevier, Amsterdam, 941 pp.
- 2247 Perry, C. L. & Mendelssohn, I. A. (2009). Ecosystem Effects of Expanding Populations of *Avicennia Germinans*
2248 in a Louisiana Salt Marsh. *Wetlands*, 29: 396–406.
- 2249 Pierce, S.M. (1983) Estimation of the non-seasonal production of *Spartina maritima* (Curtis) Fernald in a South
2250 African estuary. *Estuarine Coastal and Shelf Science* 16, 241-254.
- 2251 Phillips, R. C., McMillan, C., & Bridges, K. W. (1983). Phenology of eelgrass, *Zostera marina* L., along
2252 latitudinal gradients in North America. *Aquatic Botany*, 15(2), 145-156.
- 2253 Pillay, D., Branch, G. M., Griffiths, C. L., Williams, C., & Prinsloo, A. (2010). Ecosystem change in a South
2254 African marine reserve(1960-2009): role of seagrass loss and anthropogenic disturbance. *Marine Ecology*
2255 *Progress Series*, 415, 35-48.
- 2256 Plus, M., Chapelle, A., Ménesguen, A., Deslous-Paoli, J. M., & Auby, I. (2003). Modelling seasonal dynamics of
2257 biomasses and nitrogen contents in a seagrass meadow (*Zostera noltii* Hornem.): application to the Thau
2258 lagoon (French Mediterranean coast). *Ecological Modelling*, 161(3), 213-238.
- 2259 Plus, M., Deslous-Paoli, J. M., Auby, I., & Dagault, F. (2001). Factors influencing primary production of
2260 seagrass beds (*Zostera noltii* Hornem.) in the Thau lagoon (French Mediterranean coast). *Journal of*
2261 *experimental marine biology and ecology*, 259(1), 63-84.
- 2262 Poffenbarger, H.J, Needelman, B.A, and Megonigal, J.P. (2011). Salinity influence on methane emissions from
2263 tidal marshes. *Wetlands* 31(5):831-842.
- 2264 Poovachiranon, S., & Chansang, H. (1994). Community structure and biomass of seagrass beds in the Andaman
2265 Sea. I. Mangrove-associated seagrass beds. *Phuket Marine Biological Center Research Bulletin*, 59, 53-64.
- 2266 Pongparn, S., 2003. Common allometric relationships for estimating the biomass of mangrove forests. Ph.D.
2267 dissertation, Gifu University, 87 pp.

- 2268 Povidisa, K., Delefosse, M., & Holmer, M. (2009). The formation of iron plaques on roots and rhizomes of the
2269 seagrass *Cymodocea serrulata* (R. Brown) Ascherson with implications for sulphide intrusion. *Aquatic*
2270 *Botany*, 90(4), 303-308.
- 2271 Powell, G. V., Kenworthy, J. W., & Fourqurean, J. W. (1989). Experimental evidence for nutrient limitation of
2272 seagrass growth in a tropical estuary with restricted circulation. *Bulletin of Marine Science*, 44(1), 324-340.
- 2273 Pozo, J. and Colino, R. 1992. Decomposition processes of *Spartina maritima* in a salt marsh of the Basque
2274 Country. *Hydrobiologia* 231:165-175.
- 2275 Prasad, M.B., A.L. Ramanathan. 2009. Organic matter characterization in a tropical estuarine-mangrove
2276 ecosystem of India: preliminary assessment by using stable isotopes and lignin phenols. *ECSS* 84: 617-624.
- 2277 Preen, A. (1995). Impacts of dugong foraging on seagrass habitats: observational and experimental evidence for
2278 cultivation grazing. *Marine ecology progress series*. Oldendorf, 124(1), 201-213.
- 2279 Purvaja R. and Ramesh R. 2001. Natural and anthropogenic methane emission from coastal wetlands of South
2280 India. *Environmental Management* 27: 547-557.
- 2281 Putz, F. E., Chan, H. T. (1986). Tree growth, dynamics, and productivity in a mature mangrove forest in
2282 malaysia. *Forest ecology and management*(17), 211-230.
- 2283 Ramos e Silva, CA, Oliveira SR, Rêgo RDP, Mozeto AA. 2007. Dynamics of phosphorus and nitrogen through
2284 litter fall and decomposition in a tropical mangrove forest. *Marine Environmental Research* 64: 425-534.
- 2285 Ranjan, R. K., Routh, J., Ramanathan, A. L., & Klump, J. V. 2011. Elemental and stable isotope records of
2286 organic matter input and its fate in the Pichavaram mangrove-estuarine sediments (Tamil Nadu, India).
2287 *Marine Chemistry*, 126(1), 163-172.
- 2288 Rasheed, M. A. (1999). Recovery of experimentally created gaps within a tropical *Zostera capricorni* (Aschers.)
2289 seagrass meadow, Queensland Australia. *Journal of Experimental Marine Biology and Ecology*, 235(2), 183-
2290 200.
- 2291 Ray, R., Ganguly, D., Chowdhury, C. (2011). Carbon sequestration and annual increase of carbon stock in a
2292 mangrove forest. *Atmos. Environment*, 45:5016-5024.
- 2293 Reddy, KR and DeLaune, RD. 2008. *Biogeochemistry of Wetlands: Science and Management*. CRC Press, 774
2294 pp.
- 2295 Ren H., Chen H., Li Z.A., Han W.D. 2009. Biomass accumulation and carbon storage of four different aged
2296 *Sonneratia apetala* plantations in Southern China. *Plant and Soil* 327:279-291.
- 2297 Reyes, J., Sansón, M., & Afonso-Carrillo, J. (1995). Distribution and reproductive phenology of the seagrass
2298 *Cymodocea nodosa* (Ucria) Ascherson in the Canary Islands. *Aquatic Botany*, 50(2), 171-180.
- 2299 Richards, T. M., Krebs, J. M., & McIvor, C. C. (2011). Microhabitat associations of a semi-terrestrial fish,
2300 *Kryptolebias marmoratus* (Poey 1880) in a mosquito-ditched mangrove forest, west-central Florida. *Journal*
2301 *of Experimental Marine Biology and Ecology*, 401(1), 48-56.
- 2302 Rismondo, A., Curiel, D., Marzocchi, M., & Scattonin, M. (1997). Seasonal pattern of *Cymodocea nodosa*
2303 biomass and production in the lagoon of Venice. *Aquatic botany*, 58(1), 55-64.
- 2304 Robertson, A. I., & Phillips, M. J. 1995. Mangroves as filters of shrimp pond effluent: predictions and
2305 biogeochemical research needs. *Hydrobiologia*, 295(1), 311-321.
- 2306 Robertson, A. I. & Mann, K. H. (1984). Disturbance by ice and life-history adaptations of the seagrass *Zostera*
2307 *marina*. *Marine Biology* 80, 131-141.
- 2308 Robertson A.I. and Daniel P.A. 1989. Decomposition and the annual flux of detritus from fallen timber in
2309 tropical mangrove forests. *Limnology and Oceanography* 34: 640-646.
- 2310 Rojstaczer, S., and S. J. Deverel (1993), Time dependence in atmospheric carbon inputs from drainage of
2311 organic soils, *Geophys. Res. Lett.*, 20(13), 1383– 1386.
- 2312 Roman, C. T., & Able, K. W. (1988). Production ecology of eelgrass (*Zostera marina* L.) in a cape cod salt
2313 marsh-estuarine system, Massachusetts. *Aquatic Botany*, 32(4), 353-363.

Second Order Draft

- 2314 Roman, C.T., Daiber, F.C. (1984) Aboveground and belowground primary production dynamics of two
2315 Delaware Bay tidal marshes. *Bulletin of the Torrey Botanical Society* 111, 34-41.
- 2316 Roman, C. T., Peck, J. A., Allen, J. R., King, J. W., & Appleby, P. G. (1997). Accretion of a New England (USA)
2317 salt marsh in response to inlet migration, storms, and sea-level rise. *Estuarine, Coastal and Shelf Science*,
2318 45(6), 717-727.
- 2319 Ross, M., Ruiz, P., Telesnicki, G., Meeder, J. (2001). Estimating above-ground biomass and production in
2320 mangrove communities of Biscayne National Park, Florida (U.S.A). *Wetland Ecology and Management* (9:),
2321 27-37.
- 2322 Rublee, P. A., & Roman, M. R. (1982). Decomposition of turtlegrass (*Thalassia testudinum* Konig) in flowing
2323 sea-water tanks and litterbags: Compositional changes and comparison with natural particulate matter.
2324 *Journal of Experimental Marine Biology and Ecology*, 58(1), 47-58.
- 2325 Saenger, P. 2002. Mangrove ecology, silviculture, and conservation. Kluwer, Dordrecht.
- 2326 Saenger, P. and Snedaker, S. C. 1993. Pantropical trends in mangrove aboveground biomass and annual litterfall.
2327 *Oecologia* 96: 293-299.
- 2328 Saintilan, N. 1997. Above-and below-ground biomasses of two species of mangrove on the Hawkesbury River
2329 estuary, New South Wales. *Marine and freshwater research*, 48(2), 147-152.
- 2330 Sand-Jensen, K., & Borum, J. (1983). Regulation of growth of eelgrass (*Zostera marina* L.) in Danish coastal
2331 waters. *Marine Technology Society Journal*, 17(2), 15-21.
- 2332 Schwarz, A. M., Morrison, M., Hawes, I., & Halliday, J. (2006). Physical and biological characteristics of a rare
2333 marine habitat: sub-tidal seagrass beds of offshore islands. Science & Technical Pub., Department of
2334 Conservation.
- 2335 Scarton, F., Day, J.W., Rismondo, A. (2002) Primary production and decomposition of *Sarcocornia fruticosa* (L.)
2336 Scott and *Phragmites australis* Trin. ex Steudel in the Po Delta, Italy. *Estuaries* 25, 325-336.
- 2337 Schubauer, J.P., Hopkinson, C.S. (1984) Above- and belowground emergent macrophyte production and
2338 turnover in a coastal marsh ecosystem, Georgia. *Limnology and Oceanography* 29, 1052-1065.
- 2339 Seitzinger, S.P. and Harrison, J.A. (2008). Land-based nitrogen sources and their delivery to coastal systems.
2340 Chapter 9, pp. 469-510. In: C. Capone, D.A. Bronk, M.R. Mullholland, E. Carpenter (editors), *Nitrogen in*
2341 *the Marine Environment*. 2nd edition, Elsevier.
- 2342 Seitzinger, S. P.; Kroeze, C.; Styles, R. V. 2000. Global distribution of N₂O emissions from aquatic systems:
2343 natural emissions and anthropogenic effects *Chemosphere Global Change Sci.* 2(3-4) 267- 279
- 2344 Setia, R., Marschner, P., Baldock, J., Chittleborough, D. and Verma, V. 2011. Relationships between carbon
2345 dioxide emission and soil properties in salt-affected landscapes. *Soil Biology and Biochemistry* 43:667-674.
- 2346 Sfriso A., Ghetti P.F. (1998) Seasonal variation in biomass, morphometric parameters and production of
2347 seagrasses in the lagoon of Venice. *Aquatic Botany*, 61, 207-223.
- 2348 Sherman, R., Fahey, T., Martinez, P. (2003). Spatial of Biomass and above-ground net primary productivity in a
2349 mangrove ecosystem in the Dominican Republic. *Ecosystems* (6:), 384-398.
- 2350 Sheppard, C., & Price, A. 1991. Will marine life survive the Gulf War? *New Scientist*, 129(1759), 36-40.
- 2351 Sifleet, S., Pendleton, L., Murray, B. C. 2011. State of the Science on Coastal Blue Carbon: A summary for
2352 policy makers. Nicholas Institute for Environmental Policy Solutions Report NIR 11-06, May 2011.
- 2353 Silva CAR, Mozeto AA, Ovalle ARC. 1998. Distribution and fluxes as macrodetritus of phosphorus in red
2354 mangroves, Sepetiba Bay, Brazil. *Mangroves and Salt Marshes* 2:37-42.
- 2355 Simard, M., Zhang, K., Rivera-Monroy, V. H., Ross, M. S., Ruiz, P. L. Castañeda-Moya, Twilley, R. R.,
2356 Rodriguez, E. 2006. Mapping height and biomass of mangrove forests in Everglades National Park with
2357 SRTM elevation data. *Photogrammetric Engineering and Remote Sensing* 72: 299-311.
- 2358 Simpson LT, IC Feller, SK Chapman. 2012. Effects of competition and nutrient enrichment on *Avicennia*
2359 *germinans* in a salt marsh-mangrove ecotone. *Aquatic Botany* 104: 55-59.

- 2360 Slim, F., Gwada, P., Kodjo, M., Hemminga, M. (1996). Biomass and litter fall of *Ceriops tagal* and *Rhizophora*
2361 *mucronata* in the mangrove forest of Gazi Bbay, Kenya. Marine and. Freshwater Research .(47:), 999-1007.
- 2362 Smith, K.K., Good, R.E., Good, N.F. (1978) Production dynamics for above and belowground components of a
2363 New Jersey *Spartina alterniflora* tidal marsh. Estuarine and Coastal Marine Science 9, 189-201.
- 2364 Smith, T. J. and Whelan, K. R. T. 2006. Development of allometric relations for three mangrove species in South
2365 Florida for use in the Greater Everglades Ecosystem restoration. Wetlands Ecology and Management 14:
2366 409-419.
- 2367 Soares, M. L. G. and Schaeffer-Novelli, Y. 2005. Above-ground biomass of mangrove species. I. Analysis of
2368 models. Estuarine, Coastal and Shelf Science 65: 1-18.
- 2369 Sousa, A.I., Lillebø, A.I., Pardal, M.A., Caçador, I. (2010) Productivity and nutrient cycling in salt marshes:
2370 Contribution to ecosystem health. Estuarine, Coastal and Shelf Science 87, 640-646.
- 2371 Spain, A. V. and Holt, J. A. 1980. The elemental status of the foliage and branchwood of seven mangrove
2372 species from northern Queensland, Division of Soils, Divisional Report No49. CSIRO, Canberra.
- 2373 Squiers, E.R., Good, R.E. (1974) Seasonal changes in the productivity, caloric content, and chemical
2374 composition of a population of salt-marsh cord-grass (*Spartina alterniflora*). Chesapeake Science 15, 63-71.
- 2375 Steinke, T., Ward, C., Rajh, A. (1995). Forest structure and biomass of mangroves in the Mgeni estuary South
2376 Africa. Hydrobiologia (295:), 159-166.
- 2377 Steward, J. S., and W. Green. 2007. Setting load limits for nutrients and suspended solids based upon seagrass
2378 depthlimit targets. Estuar. Coasts 30: 657-670.
- 2379 Sukardjo, S. 1993. The present status of the mangrove forests in the northern coast of West Java with special
2380 reference to the recent utilization. Southeast Asian Studies 31: 141-157.
- 2381 Suzuki, E., Tagawa, E. (1983). Biomass of a mangrove forest and a sedge marsh on Ishigaki island, South
2382 Japan. Japan Journal Ecology (33:), 231-234.
- 2383 Tam, N. F. Y., Wong, Y. S., Lan, C. Y., & Chen, G. Z. 1995. Community structure and standing crop biomass of
2384 a mangrove forest in Futian Nature Reserve, Shenzhen, China. Hydrobiologia, 295(1), 193-201.
- 2385 Tamai, S., Tabuchi, R., Ogino, K., & Nakasuga, T. 1986. Standing biomass of mangrove forests in southern
2386 Thailand. Journal of the Japanese Forestry Society, 68.
- 2387 Terrados, J., & Ros, J. D. (1992). Growth and primary production of *Cymodocea nodosa* (Ucria) Ascherson in a
2388 Mediterranean coastal lagoon: the Mar Menor (SE Spain). Aquatic botany, 43(1), 63-74.
- 2389 Terrados, J., Duarte, C. M., Fortes, M. D., Borum, J., Agawin, N. S., Bach, S., Thampanya U., Kamp-Nielsen, L.,
2390 Kenworthy, W. J., Geertz-Hansen, O. & Vermaat, J. (1998). Changes in community structure and biomass of
2391 seagrass communities along gradients of siltation in SE Asia. Estuarine, Coastal and Shelf Science, 46(5),
2392 757-768.
- 2393 Terrados, J., Grau-Castella, M., Piñol-Santiñà, D., & Riera-Fernández, P. (2006). Biomass and primary
2394 production of a 8–11m depth meadow versus < 3m depth meadows of the seagrass *Cymodocea nodosa*
2395 (*Ucria*) Ascherson. Aquatic botany, 84(4), 324-332.
- 2396 Thant, Y. M. and Kanzaki, M. 2011. Biomass and carbon sequestration in community mangrove plantations and
2397 a natural regeneration stand in the Ayeyarwady Delta, Myanmar. American Geophysical Union, Fall Meeting,
2398 abstract #B21C-0265.
- 2399 Thayer, G. W., Engel, D. W., & LaCroix, M. W. (1977). Seasonal distribution and changes in the nutritive
2400 quality of living, dead and detrital fractions of *Zostera marina*. L. Journal of Experimental Marine Biology
2401 and Ecology, 30(2), 109-127.
- 2402 Thiery A. and Puente L. 2002. Crustacean assemblage and environmental characteristics of a man-made solar
2403 saltwork in southern France, with emphasis on anostracan (Branchiopoda) population dynamics.
2404 Hydrobiologia 486: 191-200.
- 2405 Tomasko, D. A. and Lapointe, B. E. 1991. Productivity and biomass of *Thalassia testudinum* as related to water
2406 column nutrient availability and epiphyte levels: field observations and experimental studies. Marine Ecology
2407 Progress Series 75: 9-16.

Second Order Draft

- 2408 Tong, C, Wang, WQ, Zeng, CS, Marrs, R. 2010. Methane (CH₄(4)) emission from a tidal marsh in the Min River
2409 estuary, southeast China. *Journal of Environmental Science and Health Part a-Toxic/Hazardous Substances &*
2410 *Environmental Engineering* 45: 506-516.
- 2411 Townsend, E. C., & Fonseca, M. S. (1998). Bioturbation as a potential mechanism influencing spatial
2412 heterogeneity of North Carolina seagrass beds. *Marine Ecology Progress Series*, 169, 123-132.
- 2413 Twilley, R. R., Chen, R. H., Hargis, T. Carbon sinks in mangroves and their implications to carbon budget of
2414 tropical coastal ecosystems. *Water, Air and Soil Pollution* 64: 265-288.
- 2415 Twilley R.W., Lugo A.E., Patterson-Zucca C. 1986. Litter Production and Turnover in Basin Mangrove Forests
2416 in Southwest Florida. *Ecology* 67: 670-683.
- 2417 Tyler, G. (1971) Distribution and turnover of organic matter and minerals in a shore meadow ecosystem. *Studies*
2418 *in the ecology of Baltic sea-shore meadows IV. Oikos* 22, 265-291.
- 2419 Udy JW, Dennison WC (1997) Growth and physiological responses of three seagrass species to evaluated
2420 sediment nutrients in Moreton Bay, Australia. *J Exp Mar Biol Ecol* 217:253–277.
- 2421 Uku, J., & Björk, M. (2005). Productivity aspects of three tropical seagrass species in areas of different nutrient
2422 levels in Kenya. *Estuarine, Coastal and Shelf Science*, 63(3), 407-420.
- 2423 Utrera-López, M. E., & Moreno-Casasola, P. 2008. Mangrove litter dynamics in la Mancha lagoon, Veracruz,
2424 México. *Wetlands Ecology and Management*, 16(1), 11-22.
- 2425 Valiela, I. and J. M. Teal. 1974. Nutrient limitation in salt marsh vegetation, p. 547–563. In R. J. Reimold and W.
2426 H. Queen (eds.). *Ecology of Halophytes*. Academic Press, New York.
- 2427 Valiela, J.M., Teal, J.M., and Persson, N.Y. 1976. Production and dynamics of experimentally enriched salt
2428 marsh vegetation: Belowground biomass. *Limnology and Oceanography* 21:245-252.
- 2429 van den Heuvel, R.N., Hefting, M.M., Tan, N.C.G., Jetten, M.S.M., Verhoeven, J.T.A. (2009). N₂O emission
2430 hotspots at different spatial scales and governing factors for small scale hotspots. *Science of the Total*
2431 *Environment*, 407: 2325-2332.
- 2432 van Houte-Howes, K. S., Turner, S. J., & Pilditch, C. A. (2004). Spatial differences in macroinvertebrate
2433 communities in intertidal seagrass habitats and unvegetated sediment in three New Zealand estuaries.
2434 *Estuaries and Coasts*, 27(6), 945-957.
- 2435 Van Lent, F., Nienhuis, P. H., & Verschuure, J. M. (1991). Production and biomass of the seagrasses *Zostera*
2436 *noltii* Hornem. and *Cymodocea nodosa* (Ucria) Aschers. at the Banc d'Arguin (Mauritania, NW Africa): a
2437 preliminary approach. *Aquatic botany*, 41(4), 353-367.
- 2438 van Tussenbroek, B. I. (1998). Above-and below-ground biomass and production by *Thalassia testudinum* in a
2439 tropical reef lagoon. *Aquatic botany*, 61(1), 69-82.
- 2440 Vaudrey, JMP, Kemer, JN, Branco, BF, Short FT. 2010. Eelgrass recovery after nutrient enrichment reversal.
2441 *Aquatic Botany* 93: 237-243.
- 2442 Vegas-Vilarrúbia, T., Baritto, F., López, P., Meleán, G., Ponce, M. E., Mora, L., & Gómez, O. (2010). Tropical
2443 Histosols of the lower Orinoco Delta, features and preliminary quantification of their carbon storage.
2444 *Geoderma*, 155(3), 280-288.
- 2445 Vermaat, J. E., & Verhagen, F. C. (1996). Seasonal variation in the intertidal seagrass *Zostera noltii* Hornem.:
2446 coupling demographic and physiological patterns. *Aquatic Botany*, 52(4), 259-281.
- 2447 Vermaat, J. E., Agawin, N. S. R., Duarte, C. M., Fortes, M. D., Marba, N., & Uri, J. S. (1995). Meadow
2448 maintenance, growth and productivity of a mixed Philippine seagrass bed. *Marine ecology progress series*.
2449 *Oldendorf*, 124(1), 215-225.
- 2450 Vermaat, J. E., Beijer, J. A. J., Gijlstra, R., Hootsmans, M. J. M., Philippart, C. J. M., Brink, V. D. N., &
2451 Vierssen, W. V. (1993). Leaf dynamics and standing stocks of intertidal *Zostera noltii* Hornem. and
2452 *Cymodocea nodosa* (Ucria) Ascherson on the Banc d'Arguin (Mauritania). *Hydrobiologia*, 258(1), 59-72
- 2453 Vermaat, J. E., Hootsmans, M. J. M., & Nienhuis, P. H. (1987). Seasonal dynamics and leaf growth of *Zostera*
2454 *noltii* hornem., A perennial intertidal seagrass. *Aquatic botany*, 28(3), 287-299.

- 2455 Vinther, H. F., & Holmer, M. (2008). Experimental test of biodeposition and ammonium excretion from blue
2456 mussels (*Mytilus edulis*) on eelgrass (*Zostera marina*) performance. *Journal of Experimental Marine Biology*
2457 *and Ecology*, 364(2), 72-79.
- 2458 Wahbeh, M. I. (1988). Seasonal distribution and variation in the nutritional quality of different fractions of two
2459 seagrass species from Aqaba (Red Sea), Jordan. *Aquatic botany*, 32(4), 383-392.
- 2460 Wahlen, S.C. (2005) Biogeochemistry of methane exchange between natural wetlands and the atmosphere.
2461 *Environmental Engineering Science*, 22, 73-94.
- 2462 Walker, D. I. (1985). Correlations between salinity and growth of the seagrass *Amphibolis antarctica* (Labill.)
2463 Sonder & Aschers., In Shark Bay, Western Australia, using a new method for measuring production rate.
2464 *Aquatic botany*, 23(1), 13-26.
- 2465 Walker, D. I., & McComb, A. J. (1988). Seasonal variation in the production, biomass and nutrient status of
2466 *Amphibolis antarctica* (Labill.) Sonder ex Aschers. and *Posidonia australis* hook. f. in Shark Bay, Western
2467 Australia. *Aquatic botany*, 31(3), 259-275.
- 2468 Walker, D. I., Kendrick, G. A., & McComb, A. J. (1988). The distribution of seagrass species in Shark Bay,
2469 Western Australia, with notes on their ecology. *Aquatic botany*, 30(4), 305-317.
- 2470 Wallentinus, H.-G. (1973) Above-ground primary production of a *Juncetum gerardi* on a Baltic sea-shore
2471 meadow. *Oikos* 24, 200-219.
- 2472 Walters, B.B., Ronnback, P., Kovacs, J.M., Crona, B., Hussain, SA, Badola, R., Primavera, J.H., Barbier, E.,
2473 Dahdouh-Guebas, F. 2008. Ethnobiology, socio-economics and management of mangrove forests: A review.
2474 *Aquatic Botany* 89: 220-236.
- 2475 Wang, J. K. 1990. Managing shrimp pond water to reduce discharge problems. *Aquacultural engineering*, 9(1),
2476 61-73.
- 2477 Waycott, M., Duarte, C.M., Carruthers, T.J.B., Orth, R.J. and others. (2009.) Accelerating loss of seagrasses
2478 across the globe threatens coastal ecosystems. *Proc Natl Acad Sci USA* 106:12377–12381
- 2479 West, R. J., & Larkum, A. W. D. (1979). Leaf productivity of the seagrass, *Posidonia australis*, in eastern
2480 Australian waters. *Aquatic Botany*, 7, 57-65.
- 2481 Whigham D.F, McCormick J., Good R.E., and Simpson R.L. 1978. Biomass and primary production in
2482 freshwater tidal wetlands of the Middle Atlantic Coast. In *Freshwater Wetlands: Ecological Processes and*
2483 *Management Potential* (R.E. Good, D.F. Whigham, and R.L. Simpson, eds.). pp. 3-20. Academic Press, NY,
2484 NY.
- 2485 White, D.A., Weiss, T.E., Trapani, J.M., Thien, L.B. (1978) Productivity and decomposition of the dominant salt
2486 marsh plants in Louisiana. *Ecology* 59, 751-759.
- 2487 Williams, S. L. (1987). Competition between the seagrasses *Thalassia testudinum* and *Syringodium filiforme* in a
2488 Caribbean lagoon. *Mar Ecol. Prog. Ser.*, 35, 91-98.
- 2489 Williams, R.B., Murdoch, M.B. (1972) Compartmental analysis of the production of *Juncus roemerianus* in a
2490 North Carolina salt marsh. *Chesapeake Science* 13, 69-79.
- 2491 Woodroffe, C. D., & Moss, T. J. (1984). Litter fall beneath *Rhizophora stylosa* griff., Vaitupu, Tuvalu, South
2492 Pacific. *Aquatic botany*, 18(3), 249-255.
- 2493 World Bank. 2006. Aquaculture: Changing the Face of the Waters Meeting the Promise and Challenge of
2494 Sustainable Aquaculture, Report No. 36622 – GLB.
2495 http://siteresources.worldbank.org/INTARD/Resources/Aquaculture_ESW_vGDP.pdf
- 2496 Yarbro, L. A., & Carlson, P. R. (2008). Community oxygen and nutrient fluxes in seagrass beds of Florida Bay,
2497 USA. *Estuaries and coasts*, 31(5), 877-897.
- 2498 Yang D, and Yang C. 2009 Detection of seagrass distribution changes from 1991 to 2006 in Xincun Bay,
2499 Hainan, with satellite remote sensing. *Sensors* 2: 830-844.
- 2500 Yu, O. T., & Chmura, G. L. (2009). Soil carbon may be maintained under grazing in a St Lawrence Estuary tidal
2501 marsh. *Environmental Conservation*, 36(04), 312-320.

Second Order Draft

- Zainal A.J.M., Dalby D.H. and Robinson, I.S., 1993. Monitoring marine ecological changes on the east coast of Bahrain with LANDSAT TM. *Photogrammetric Engineering and Remote Sensing*, 59, 415–421.
- Zavodnik, N., Travizi, A., & De Rosa, S. (1998). Seasonal variations in the rate of photosynthetic activity and chemical composition of the seagrass *Cymodocea nodosa* (Ucr.) Asch. *Scientia Marina*, 62(4), 301-309.
- Zieman, J., Fourqurean, J. W., & Iverson, R. L. (1989). Distribution, abundance and productivity of seagrasses and macroalgae in Florida Bay. *Bulletin of Marine Science*, 44(1), 292-311.
- Zhang, J.P., Shen, C-D., Ren, H., Wang, J., Han, W-D. 2012. Estimating change in sedimentary organic carbon content during mangrove restoration in southern China using carbon isotopic measurements. *Pedosphere* 22:58-66.
- Zhang, Y., Huang, G., Wang, W., Chen, L., and Lin, G., (2012). Interactions between mangroves and exotic *Spartina* in an anthropogenically disturbed estuary in southern China, *Ecology*, **93**(3), 588–597.

ANNEX 4.1 PERCENT REFRACTORY CARBON

Percent refractory carbon in organic/mineral soils were estimated for mangrove soils based on either the amount of phenolic compounds/lignins in soils or % TOC in mangrove soils deeper than 1 m if there was no further decline in TOC concentration (Table 4.3).

PERCENT REFRACTORY CARBON APPLIED TO ESTIMATE % C OXIDATION FOR MANGROVE SOILS (% BY SOIL DRY WEIGHT)		2520
Mean	3.98	2521
Median	3.4	2522
N	16	2523
Prasad & Ramanathan 2009; Marchand et al. 2003; Dittmar & Lara 2001; Koch et al. 2011; Ranjan et al., 2010; Marchand et al. 2005), which is similar to that in tidal marshes (Filipet al. 1988; Alberts et al., 1988; Ramesh et al. 2008)		2524

ANNEX 4.2 ADDITIONAL SOURCES FOR ACTIVITY DATA

The following web sources provide links to domestic and international data sources pertaining to coastal wetland areas and other activity data for management activities contained in this chapter. Other web sources are provided in the text of the chapter in the respective activity data sections. Resources providing recent trends in coastal wetland area can help countries understand circumstances of those trends and what management activities contribute to them (FAO 2007; Green and Short 2003; Sifleet et al. 2011, <http://archive.org/stream/worldatlasofseag03gree#page/n5/mode/2up>; <http://nicholasinstitute.duke.edu/publications/>; Fatoyinbo & Simard 2013)

Mangroves

- Mangrove.org: <http://mangrove.org/>
- Mangrove Action Project: <http://www.mangroveactionproject.org/>
- Mangrove, National Geographic Magazine: <http://ngm.nationalgeographic.com/2007/02/mangroves/warne-text>
- FAO Mangrove Management: <http://www.fao.org/forestry/mangrove/en/>
- USGS National Wetlands Research Center: <http://www.nwrc.usgs.gov/index.html>
- World Atlas of Mangrove: <http://www.fao.org/forestry/20067/en/>

- 2544 World Distribution of Coral Reefs and Mangroves: <http://www.unep->
 2545 [wcmc.org/marine/data/coral_mangrove/marine.maps.main.html](http://www.unep-wcmc.org/marine/data/coral_mangrove/marine.maps.main.html)
- 2546 International Society for Mangrove Ecosystems: <http://www.mangrove.or.jp/>
- 2547 Global Mangrove Database & Information System: <http://www.glomis.com/>
- 2548 The UNESCO Mangrove Programme: <http://www.unesco.org/csi/intro/mangrove.htm>
- 2549 Mangrove and the Ramsar Convention: http://www.ramsar.org/types_mangroves.htm
- 2550 USGS Global Mangrove Project <http://lca.usgs.gov/lca/globalmangrove/index.php>
- 2551 **Mangroves, tidal marshes and seagrass meadows**
- 2552 <http://data.unep-wcmc.org/>
- 2553 Global distribution of seagrasses (V2.0, 2005) prepared by UNEP World Conservation Monitoring Centre
 2554 (UNEP-WCMC) in collaboration with Dr. Frederick T. Short.
- 2555 Global distribution of Mangroves (V3.0, 1997) compiled by UNEP World Conservation Monitoring Centre
 2556 (UNEP-WCMC) in collaboration with the International Society for Mangrove Ecosystems (ISME).

2557

2558

2559 **ANNEX 4.3 ACCURATE ESTIMATION OF ABOVEGROUND MANGROVE** 2560 **BIOMASS: HIGHER TIER METHODOLOGY**

2561 Because of field conditions and heavy weight of wood, an accurate survey of a mangrove forest is difficult and
 2562 time-consuming. Allometric methods (Soares and Schaeffer-Novelli, 2005; Komiyama et al., 2008) estimate the
 2563 whole or partial weight of a tree from measurable tree dimensions, notably trunk diameter and height, using
 2564 allometric relations developed from empirical measurement of weight of individual tree components (leaves,
 2565 branches, stem). Use of allometric equations is favored because it is non-destructive and is therefore useful for
 2566 estimating temporal changes in forest biomass by means of subsequent stem diameter measurements over
 2567 subsequent years.

2568 Up until recently, the major drawback of this method has been the site- and species-specific differences in
 2569 allometric relations, necessitating the use of different allometric equations for different sites (e.g., Smith and
 2570 Whelan, 2005) and, at a minimum, different species. However, a number of workers, using global datasets, have
 2571 developed a common allometric equation applicable for all tropical tree species, with the most applicable
 2572 equations for aboveground biomass being those developed for all tropical trees by Chave et al. (2005) and for all
 2573 mangrove species by Komiyama et al. (2005):

$$2574 \quad W_{top} = 0.168pDBH^{2.47} \text{ (Chave et al. 2005)}$$

$$2575 \quad W_{top} = 0.251pD^{2.46} \text{ (Komiyama et al. 2005)}$$

2576 where W_{top} = aboveground tree weight in kg DW; D = tree diameter; DBH = diameter-at-breast height. The
 2577 relative error of each equation varies among species, but is typically within the range of -10% to +10%. There
 2578 are, of course, arguments to be made that empirical measurements should be made in all mangrove forests,
 2579 considering the significant allometric differences between species and for the same species at different locations
 2580 (Smith and Whelan, 2005; Soares and Schaeffer-Novelli, 2005). However, this idea is impractical for inventory
 2581 compliers; a relative error of $\pm 10\%$ is acceptable being within the range of error for allometric relations within a
 2582 forest where biomass has been weighted.

2583 Comparing the two equations, the Chave estimation gives lower aboveground weight estimates than that of the
 2584 Komiyama equation. Presuming that a complete census of all trees, with species identified, and their diameter
 2585 have been undertaken from replicate plots within a given forest, these numbers can then be used in either
 2586 equation to derive individual tree weight.